

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 November 2004 (04.11.2004)

PCT

(10) International Publication Number
WO 2004/094659 A1

(51) International Patent Classification⁷: C12Q 1/68

PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(21) International Application Number:
PCT/EP2004/004178

(22) International Filing Date: 20 April 2004 (20.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/464,372 21 April 2003 (21.04.2003) US
60/526,689 2 December 2003 (02.12.2003) US

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for the following design-
ations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW,
BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC,
EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ,
OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,
SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
BG, CH, CY, CZ, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT,
LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF,
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
TD, TG)
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for the following design-
ation DE
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(71) Applicant (for DE only): **ROCHE DIAGNOSTICS**
GMBH [DE/DE]; Sandhofer Strasse 116, 68305
Mannheim (DE).

(71) Applicant (for all designated States except DE): **F. HOFF-
MANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse
124, CH-4070 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HIGUCHI, Russell,**
Gene [US/US]; 3258 Liberty Avenue, Alameda, CA 94501
(US). **PELTZ, Gary, Allen** [US/US]; 110 Danbury Lane,
Redwood City, CA 94061 (US). **FLJAL, Bonnie** [US/US];
59 Taurus Drive #3B, Hillsborough, NJ 08844 (US). **RO,**
Sunhee, Kwon [KR/US]; 635 Greenwich Lane, Foster
City, CA 94404 (US). **LI, Jia** [US/US]; 32801 Regents
Boulevard, Union City, CA 94587 (US).

(74) Common Representative: **ROCHE DIAGNOSTICS**
GMBH; Patent Department, 68298 Mannheim (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,

(54) Title: ASSOCIATIONS OF POLYMORPHISMS IN THE FRZB GENE WITH OBESITY AND OSTEOPOROSIS

(57) Abstract: Methods and reagents for determining sequence variants present at the FRZB locus, which facilitate identifying individuals at risk for obesity and/or osteoporosis, are provided. In particular, methods for detecting an individual's risk for obesity and/or osteoporosis by detecting at least one polymorphism (e.g., at least one single nucleotide polymorphism) in the FRZB gene in a nucleic acid sample from the individual are provided. Related kits, arrays and systems are also described.

WO 2004/094659 A1

ASSOCIATIONS OF POLYMORPHISMS IN THE FRZB GENE WITH
OBESITY AND OSTEOPOROSIS

5 FIELD OF THE INVENTION

The invention relates to methods and reagents for detecting an individual's risk for obesity and/or osteoporosis. More specifically, it relates to methods and reagents for detecting an individual's increased or decreased risk for obesity and/or osteoporosis by identifying the presence of at least one polymorphism in the FRZB gene.

10 BACKGROUND OF THE INVENTION

Transplantation experiments by Spemann and Mangold (1924, *Arch. Mikroskopische Anat. Entwicklungsmechanik*, 100:599-638) established the presence of an anatomically discrete region, the Spemann organizer, or dorsal lip, that controls patterning of the developing body axis in vertebrate embryos. Diffusible factors emanating from this region were found to be involved in different developmental processes. The arrangement of *Drosophila* cuticle hairs in a defined polarity was found to be genetically controlled by 'frizzled' (FZD1), a 7-transmembrane receptor with a large extracellular cysteine-rich domain.

The bovine and human homologs of FZD1 were cloned by RT-PCR and by screening bovine articular cartilage and human placenta cDNA libraries, which identified cDNAs encoding FRZB, the mammalian analog of FZD1 (Hoang *et al.*, 1996, *J. Biol. Chem.*, 271:26131-26137). The deduced 325-amino acid bovine and human FRZB proteins share 94% amino acid identity. Sequence analysis predicted that FRZB contains a 25-amino acid signal peptide, an N-terminal N-glycosylation site, a 24-amino acid putative transmembrane segment, a region with multiple potential Ser/Thr phosphorylation sites, and a serine-rich C-terminal domain. The N-terminal region of FRZB shares 50% amino acid identity, including the conservation of 10 Cys residues, with frizzled.

Immunoblot analysis determined that FRZB is expressed as an approximately 36-kD protein. In situ hybridization analysis of human embryos representing different stages of development detected no expression from week 6 through week 13 except in the developing appendicular skeleton, as well as in several craniofacial bones and epiphyseal

CONFIRMATION COPY

- 2 -

ends of the rib cage. Immunochemical analysis confirmed the expression of FRZB in the developing skeletal structures.

Northern blot analysis revealed that FRZB is expressed strongly in placenta and heart, at intermediate levels in brain, skeletal muscle, kidney, and pancreas, and at low levels in lung and liver (Leyns *et al.*, 1997, *Cell* 88:747-756). SDS-PAGE analysis detected secretion of FRZB, possibly after proteolytic cleavage, consistent with FRZB's lack of the 7 transmembrane domains found in the *Drosophila* and vertebrate frizzled gene family. Functional analysis in *Xenopus* embryos showed that FRZB can antagonize the early and late effects of WNT8 signaling. Mammalian WNT genes include oncogenes that lead to mammary tumors. For further characterization of FRZB, see, e.g., Dann *et al.*, 2001, *Nature* 412:86-90; Rattner *et al.*, 1997, *Proc. Nat. Acad. Sci.* 94:2859-2863; and Schumann *et al.*, 2000, *Cardiovasc. Res.* 45:720-728.

Leyns *et al.* (*supra*) mapped the human FRZB gene to 2q31-q33. They noted that loss of one copy of the 2q arm occurs with high incidence in lung and colorectal carcinomas, as well as in neuroblastomas, and suggested that FRZB might function as a tumor suppressor gene. Hoang *et al.* (*supra*) suggested FRZB might play a role in skeletal morphogenesis. However, a direct role for the FRZB gene in human disease and development has not been identified.

Among other aspects, the present invention provides alleles of FRZB, identified by the presence of one or more predisposing or protective polymorphisms, that are associated with an increased or decreased risk for obesity and/or osteoporosis. A complete understanding of the invention will be obtained upon review of the following.

SUMMARY OF THE INVENTION

The invention provides methods, reagents and kits for detecting an individual's increased or decreased risk for obesity and/or osteoporotic diseases and related diseases. In certain embodiments, the methods are used to determine an individual's risk for obesity and related diseases. In other embodiments, the methods are used to determine an individual's risk for osteoporosis and related diseases. In further embodiments, the methods of determining risk are combined with known clinical methods to diagnose osteoporosis or obesity.

- 3 -

A first general class of embodiments provides methods for determining an individual's risk for obesity. In the methods, presence of at least one obesity-related polymorphism in a frizzled-related protein (FRZB) gene in a nucleic acid sample of the individual is detected. The presence of the at least one polymorphism provides an indication of the individual's risk for obesity. The individual's risk for obesity can be, e.g., either an increased risk or a decreased risk as compared to an individual without the at least one polymorphism (e.g., an individual with a different allele at that polymorphic site). Accordingly, the at least one polymorphism can comprise a predisposing or a protective polymorphism in the FRZB gene.

- 10 The at least one polymorphism can comprise essentially any suitable polymorphism(s), including, but not limited to, restriction fragment length polymorphisms, random amplified polymorphic DNA, arbitrary fragment length polymorphisms, simple sequence repeats, single-stranded conformation polymorphisms, and amplified variable sequences. In a preferred class of embodiments, the at least one polymorphism
- 15 comprises at least one single nucleotide polymorphism (SNP). For example, the polymorphism can be either allele of T2303723C, C18679T, G19524A, T19575G, T22242A, G23043A, G23415A, T23549C, A24791G, C26794G, or G27014A. In one class of embodiments, the at least one polymorphism is selected from the group consisting of: T allele of T2303723C, C allele of T2303723C, C allele of C18679T, T allele of C18679T,
- 20 G allele of G19524A, A allele of G19524A, T allele of T22242A, A allele of T22242A, A allele of A24791G, and G allele of A24791G.

In one class of embodiments, the presence of two or more polymorphisms is detected (e.g., two or more polymorphisms at a single polymorphic site and/or at different polymorphic sites, e.g., a haplotype). At least one of the two or more polymorphisms

25 (e.g., all of the polymorphisms) is optionally selected from the group consisting of: the T allele of T2303723C, the C allele of T2303723C, the C allele of C18679T, the T allele of C18679T, the G allele of G19524A, the A allele of G19524A, the T allele of T22242A, the A allele of T22242A, the A allele of A24791G, and the G allele of A24791G.

The nucleic acid sample typically comprises DNA or RNA. The presence of the at least

30 one polymorphism in the nucleic acid sample can be detected by any of the variety of methods known in the art. For example, the at least one polymorphism can be detected by sequencing, e.g., sequencing of the region of the FRZB gene including the polymorphic site(s). The region of FRZB is optionally amplified prior to the sequencing step. As another example, the at least one polymorphism is detected by amplification,

e.g., of the region of the FRZB gene including the polymorphic site(s). The amplification can be, e.g., an allele-specific amplification. The amplification can comprise a polymerase chain reaction (e.g., kinetic PCR), a ligase chain reaction, or the like. As yet another example, the polymorphism can be detected by hybridization of a nucleic acid probe. Thus, in one class of embodiments, to detect the polymorphism the nucleic acid sample is contacted with at least one sequence-specific oligonucleotide under conditions (e.g., stringent conditions) that allow binding of the at least one oligonucleotide to the nucleic acid sample. The at least one sequence-specific oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the at least one obesity-related polymorphism. Hybridization of the at least one oligonucleotide to the nucleic acid sample is then detected. In a related class of embodiments, the FRZB region comprising the polymorphism(s) is amplified prior to probe hybridization. Thus, in this class of embodiments, the nucleic acid sample is amplified to provide an amplified nucleic acid sample. The amplified nucleic acid sample is contacted with at least one sequence-specific oligonucleotide under conditions that allow binding of the oligonucleotide to the amplified nucleic acid sample (e.g., stringent conditions). The at least one sequence-specific oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the at least one obesity-related polymorphism. Hybridization of the at least one sequence-specific oligonucleotide to the amplified nucleic acid sample is detected. The presence of the at least one obesity-related polymorphism can be, e.g., qualitatively or quantitatively detected.

In certain embodiments, the presence of the polymorphism inherited from one of the individual's parents provides an indication of the individual's risk for obesity (e.g., when the associated FRZB allele exerts a dominant effect, such that inheritance of the polymorphism from one parent is sufficiently predictive). In other embodiments, the presence of the polymorphism inherited from both of the individual's parents provides an indication of the individual's risk for obesity.

As noted, the methods are optionally combined with known clinical methods, e.g., to diagnose obesity. Thus, the methods optionally include performing at least one clinical test for obesity (e.g., determining a body mass index (BMI) of the individual).

A second general class of embodiments provides methods for determining an individual's risk for osteoporosis. In the methods, presence of at least one osteoporosis-related polymorphism in a frizzled-related protein (FRZB) gene in a nucleic acid sample

of the individual is detected. The presence of the at least one polymorphism provides an indication of the individual's risk for osteoporosis. The individual's risk for osteoporosis can be, e.g., either an increased risk or a decreased risk as compared to an individual without the at least one polymorphism (e.g., an individual with a different allele at that polymorphic site). Accordingly, the at least one polymorphism can
5 comprise a predisposing or a protective polymorphism in the FRZB gene.

The at least one polymorphism can comprise essentially any suitable polymorphism(s), including, but not limited to, restriction fragment length polymorphisms, random amplified polymorphic DNA, arbitrary fragment length polymorphisms, simple
10 sequence repeats, single-stranded conformation polymorphisms, and amplified variable sequences. In a preferred class of embodiments, the at least one polymorphism comprises at least one single nucleotide polymorphism (SNP). For example, the polymorphism can be either allele of T2303723C, C18679T, G19524A, T19575G, T22242A, G23043A, G23415A, T23549C, A24791G, C26794G, or G27014A. In one class
15 of embodiments, the at least one polymorphism is selected from the group consisting of: C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, A allele of A24791G, G allele of A24791G, C allele of C26794G, G allele of C26794G, G allele of G27014A, and A allele of G27014A.

In one class of embodiments, the presence of two or more polymorphisms is detected
20 (e.g., two or more polymorphisms at a single polymorphic site and/or at different polymorphic sites, e.g., a haplotype). At least one of the two or more polymorphisms (e.g., all of the polymorphisms) is optionally selected from the group consisting of: the C allele of C18679T, the T allele of C18679T, the G allele of G19524A, the A allele of G19524A, the A allele of A24791G, the G allele of A24791G, the C allele of C26794G, the
25 G allele of C26794G, the G allele of G27014A, and the A allele of G27014A.

The nucleic acid sample typically comprises DNA or RNA. The presence of the at least one polymorphism in the nucleic acid sample can be detected by any of the variety of methods known in the art. For example, the at least one polymorphism can be detected by sequencing, e.g., sequencing of the region of the FRZB gene including the
30 polymorphic site(s). The region of FRZB is optionally amplified prior to the sequencing step. As another example, the at least one polymorphism is detected by amplification, e.g., of the region of the FRZB gene including the polymorphic site(s). The amplification can be, e.g., an allele-specific amplification. The amplification can comprise a polymerase chain reaction (e.g., kinetic PCR), a ligase chain reaction, or the

- 6 -

- like. As yet another example, the polymorphism can be detected by hybridization of a nucleic acid probe. Thus, in one class of embodiments, to detect the polymorphism the nucleic acid sample is contacted with at least one sequence-specific oligonucleotide under conditions (e.g., stringent conditions) that allow binding of the at least one
- 5 oligonucleotide to the nucleic acid sample. The at least one sequence-specific oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the at least one osteoporosis-related polymorphism. Hybridization of the at least one oligonucleotide to the nucleic acid sample is then detected. In a related class of
- 10 embodiments, the FRZB region comprising the polymorphism(s) is amplified prior to probe hybridization. Thus, in this class of embodiments, the nucleic acid sample is amplified to provide an amplified nucleic acid sample. The amplified nucleic acid sample is contacted with at least one sequence-specific oligonucleotide under conditions that allow binding of the oligonucleotide to the amplified nucleic acid sample (e.g., stringent conditions). The at least one sequence-specific oligonucleotide hybridizes
- 15 under stringent conditions to a region of the FRZB gene comprising the at least one osteoporosis-related polymorphism. Hybridization of the at least one sequence-specific oligonucleotide to the amplified nucleic acid sample is detected. The presence of the at least one osteoporosis-related polymorphism can be, e.g., qualitatively or quantitatively detected.
- 20 In certain embodiments, the presence of the polymorphism inherited from one of the individual's parents provides an indication of the individual's risk for osteoporosis (e.g., when the associated FRZB allele exerts a dominant effect, such that inheritance of the polymorphism from one parent is sufficiently predictive). In other embodiments, the presence of the polymorphism inherited from both of the individual's parents provides
- 25 an indication of the individual's risk for osteoporosis.

As noted, the methods are optionally combined with known clinical methods, e.g., to diagnose osteoporosis. Thus, the methods optionally include performing at least one clinical test for osteoporosis (e.g., a bone-turnover assay or a bone scan).

- Yet another general class of embodiments provides methods for determining an
- 30 individual's risk for obesity and/or osteoporosis. In the methods, the individual's genotype at one or more polymorphic sites in an FRZB gene is determined. A first genotype at the one or more polymorphic sites is statistically associated with an increased risk for obesity and/or osteoporosis as compared to a second genotype at the one or more polymorphic sites. Thus, for example, if the individual's genotype

corresponds to the first genotype, the individual's risk for obesity and/or osteoporosis is greater than that of other individuals who have the second genotype.

Each polymorphic site can comprise one or more nucleotides. In a preferred class of embodiments, at least one of the one or more polymorphic sites consists of a single
5 nucleotide position (i.e., the individual is genotyped for one or more SNPs, e.g., a plurality and/or a haplotype of SNPs). In certain embodiments, a plurality of the polymorphic sites each consists of a single nucleotide position (e.g., position 2628, 18679, 19524, 19575, 22242, 23043, 23415, 23549, 24791, 26794, or 27014 of SEQ ID NO:1). For example, at least one of the one or more polymorphic sites can be selected
10 from the group consisting of: nucleotide position 2628, nucleotide position 18679, nucleotide position 19524, nucleotide position 22242, nucleotide position 24791, nucleotide position 26794, and nucleotide position 27014 of SEQ ID NO:1.

In some embodiments, the presence of a single allele of a particular polymorphism is sufficient to indicate whether the individual's risk of obesity and/or osteoporosis is
15 increased or decreased. In other embodiments, two copies of an allele of a particular polymorphism must be present to indicate an increased or decreased risk of obesity and/or polymorphism (e.g., when the effect is recessive such that both homologous chromosomes must carry the allele). Thus, in one class of embodiments, the first genotype is statistically associated with an increased risk for obesity as compared to the
20 second genotype. In one class of example embodiments, the first genotype comprises two T alleles and the second genotype two C alleles or one T allele and one C allele of T2303723C; the first genotype comprises two T alleles and the second genotype two C alleles or one T allele and one C allele of SNP C18679T; the first genotype comprises two A alleles and the second genotype two G alleles or one A allele and one G allele of SNP
25 G19524A; the first genotype comprises two A alleles and the second genotype two T alleles or one A allele and one T allele of SNP T22242A; and/or the first genotype comprises two G alleles and the second genotype two A alleles or one G allele and one A allele of SNP A24791G. In a related class of embodiments, the first genotype is statistically associated with an increased risk for osteoporosis as compared to the second
30 genotype. The first genotype comprises two C alleles and the second genotype two T alleles or one T allele and one C allele of SNP C18679T; the first genotype comprises two G alleles and the second genotype two A alleles or one A allele and one G allele of SNP G19524A; the first genotype comprises two A alleles and the second genotype two G alleles or one G allele and one A allele of SNP A24791G; the first genotype comprises
35 two C alleles and the second genotype two G alleles or one C allele and one G allele of

- 8 -

SNP C26794G; and/or the first genotype comprises two G alleles and the second genotype two A alleles or one G allele and one A allele of SNP G27014A.

Determining the individual's genotype typically involves obtaining a nucleic acid sample from the individual. Determining the individual's genotype can involve amplifying at
5 least a portion of the FRZB gene from the nucleic acid sample, the portion comprising at least one of the one or more polymorphic sites. Such amplification can be, e.g., to directly determine the genotype or to facilitate detection of one or more polymorphisms by an additional step. In one class of embodiments, the individual's genotype is determined by performing an allele-specific amplification or an allele-specific extension
10 reaction. In another class of embodiments, the individual's genotype is determined by sequencing at least a portion of the FRZB gene from the nucleic acid sample, the portion comprising at least one of the one or more polymorphic sites. In yet another class of embodiments, the individual's genotype is determined by hybridization of a nucleic acid probe, optionally after amplification of at least a portion of the FRZB gene. In one class
15 of example embodiments, at least one of the one or more polymorphic sites consists of a single nucleotide position. In these embodiments, the nucleic acid sample is contacted with at least one sequence-specific oligonucleotide under stringent conditions. The oligonucleotide hybridizes under the stringent conditions to the nucleic acid sample when a first nucleotide occupies the nucleotide position defining the polymorphic site
20 but not when a second nucleotide occupies the nucleotide position. Hybridization of the oligonucleotide to the nucleic acid sample is detected.

One aspect of the invention provides kits for detecting presence of a first predisposing or protective polymorphism in an FRZB gene, e.g., in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed. Thus, one general class of
25 embodiments provides a kit including one or more first oligonucleotides capable of detecting the first polymorphism and instructions for detecting the first polymorphism with the one or more first oligonucleotides and for correlating said detection to the individual's risk for osteoporosis and/or obesity, packaged in one or more containers.

Essentially all of the features noted for the method embodiments above apply to this
30 embodiment as well, as relevant. For example, in a preferred class of embodiments, the first polymorphism is a single nucleotide polymorphism, e.g., a SNP selected from the group consisting of: the T allele of T2303723C, the C allele of T2303723C, the C allele of C18679T, the T allele of C18679T, the G allele of G19524A, the A allele of G19524A, the T allele of T22242A, the A allele of T22242A, the A allele of A24791G, the G allele of

- 9 -

A24791G, the C allele of C26794G, the G allele of C26794G, the G allele of G27014A, and the A allele of G27014A. Other potential SNPs include, but are not limited to, either allele of T19575G, G23043A, G23415A, and T23549C.

In one aspect, the kit can be used to detect the presence of the first polymorphism by hybridization of a nucleic acid probe to the polymorphism. Thus, in one class of
5 embodiments, the one or more first oligonucleotides comprise at least one probe. In certain embodiments, the first oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the first polymorphism. In one class of
10 embodiments, the first polymorphism is a first single nucleotide polymorphism comprising a first nucleotide at a first nucleotide position. In this class of embodiments, under stringent conditions, the first oligonucleotide hybridizes to a region of the FRZB
15 gene comprising the first single nucleotide polymorphism with a signal to noise ratio that is at least 2x (e.g., at least 5x or at least 10x) the signal to noise ratio at which the first oligonucleotide hybridizes to the region of the FRZB gene comprising a second
20 nucleotide at the first nucleotide position. The first oligonucleotide is typically fully complementary to the region of the FRZB gene comprising the first polymorphism, and typically comprises at least about 10 contiguous nucleotides complementary to the
25 FRZB gene.

To facilitate detection of the polymorphism (e.g., through detection of hybridization
20 between the one or more first oligonucleotides and a nucleic acid comprising the polymorphism), for example, the one or more first oligonucleotides optionally comprise a label, e.g., an isotopic, fluorescent, fluorogenic, luminescent or colorimetric label. In some embodiments, the label itself directly produces a detectable signal (e.g., a
25 fluorescent label). In other embodiments, the kit also includes a reagent that detects the label (e.g., an enzyme that cleaves a colorimetric label, a binding moiety, or the like).

In one aspect, the one or more first oligonucleotides comprise one or more primers. The primer(s) can be used to detect the polymorphism, e.g., in an allele-specific
30 amplification or extension reaction. For example, in one class of embodiments, the first polymorphism is a first single nucleotide polymorphism comprising a first nucleotide at a first nucleotide position, and the 3' nucleotide of one of the one or more first
35 oligonucleotides is complementary to the first nucleotide.

The primer(s) can be used to amplify a region of FRZB comprising the polymorphism, e.g., for subsequent detection of the polymorphism by hybridization, sequencing, or the

like. In one class of embodiments, the one or more first oligonucleotides comprise amplification primers, wherein the amplification primers amplify a nucleic acid sequence comprising the first polymorphism. In a related class of embodiments, the one or more first oligonucleotides comprise sequencing primers that flank the first
5 polymorphism.

The one or more first oligonucleotides are optionally immobilized on a substrate. The substrate can be, for example, a planar substrate or a beaded substrate. The oligonucleotide(s) can be arranged in an array of other oligonucleotides used to detect other polymorphisms, e.g., other polymorphisms in FRZB.

10 The kit can optionally be used to detect more than one polymorphism (simultaneously or sequentially). Thus, in one class of embodiments, the kit also includes one or more second oligonucleotides capable of detecting a second polymorphism (and optionally third, fourth, fifth, etc. oligonucleotides capable of detecting third, fourth, fifth, etc. polymorphisms). The second polymorphism can be at the same polymorphic site as the
15 first or at a different polymorphic site (in FRZB or a different gene), and can be protective or predisposing.

One general class of embodiments provides arrays for detecting presence of one or more predisposing and/or protective polymorphisms in an FRZB gene, e.g., in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed. In
20 one class of embodiments, the array comprises a substrate and a plurality of oligonucleotides, each of which oligonucleotides hybridizes to a region of the FRZB gene comprising at least one of the polymorphisms. The hybridization detects the presence of the polymorphism, and this detection provides an indication of the individual's risk for osteoporosis and/or obesity. The plurality of oligonucleotides are immobilized on the
25 substrate. Typically, the array is used for detecting the presence of a plurality of polymorphisms, e.g., multiple alleles at a single polymorphic site and/or different polymorphic sites.

Essentially all of the features noted for the method and kit embodiments above apply to this embodiment as well, as relevant. For example, the one or more polymorphisms
30 preferably comprise one or more single nucleotide polymorphisms. For example, at least one of the one or more polymorphisms can be selected from the group consisting of: the T allele of T2303723C, the C allele of T2303723C, the C allele of C18679T, the T allele of C18679T, the G allele of G19524A, the A allele of G19524A, the T allele of

- 11 -

T22242A, the A allele of T22242A, the A allele of A24791G, the G allele of A24791G, the C allele of C26794G, the G allele of C26794G, the G allele of G27014A, and the A allele of G27014A. Other potential SNPs include, but are not limited to, either allele of T19575G, G23043A, G23415A, and T23549C.

- 5 In one class of embodiments in which the array can be used to detect presence of one or more SNPs, each of the oligonucleotides in the array hybridizes under stringent conditions to a region of the FRZB gene comprising one of the single nucleotide polymorphisms with a signal to noise ratio that is at least 2x (e.g., at least 5x or at least 10x) that at which the oligonucleotide hybridizes to a region of the FRZB gene
10 comprising any of the remaining single nucleotide polymorphisms. Typically, one oligonucleotide is used to detect one SNP; that is, each of the oligonucleotides typically hybridizes to a distinct single nucleotide polymorphism.

As noted, the plurality of oligonucleotides are immobilized on a substrate, e.g., a planar substrate, a membrane, a glass slide, or the like. Typically, each of the plurality of
15 oligonucleotides is immobilized at a known, pre-determined position on the substrate.

To facilitate detection of polymorphisms by specific hybridization with the oligonucleotides, each of the plurality of oligonucleotides is typically fully complementary to a region of the FRZB gene comprising one of the polymorphisms, and each of the plurality of oligonucleotides typically comprises at least about 10
20 contiguous nucleotides complementary to the FRZB gene. Each of the plurality of oligonucleotides optionally comprises a label, e.g., a label that facilitates detection of hybridization between the oligonucleotide and the corresponding polymorphism.

The array is optionally part of a system. Thus, one class of embodiments provides a system comprising an array of the invention and system instructions that correlate the
25 detection of the presence of one or more predisposing or protective polymorphisms to the individual's risk for osteoporosis and/or obesity.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts exemplary FRZB SNPs. Nine base pairs centered on each SNP are
30 shown; the upper strand corresponds to the strand whose sequence is listed in SEQ ID NO:1. SNP alleles illustrated are: C allele of C18679T (Panel A), T allele of C18679T (Panel B), G allele of G19524A (Panel C), A allele of G19524A (Panel D), T allele of

- 12 -

T19575G (Panel E), G allele of T19575G (Panel F), T allele of T22242A (Panel G), A allele of T22242A (Panel H), G allele of G23043A (Panel I), A allele of G23043A (Panel J), G allele of G23415A (Panel K), A allele of G23415A (Panel L), T allele of T23549C (Panel M), C allele of T23549C (Panel N), A allele of A24791G (Panel O), G allele of A24791G (Panel P), C allele of C26794G (Panel Q), G allele of C26794G (Panel R), G allele of G27014A (Panel S), A allele of G27014A (Panel T), T allele of T2303723C (Panel U), and C allele of T2303723C (Panel V).

DEFINITIONS

10 Before describing the invention in detail, it is to be understood that this invention is not limited to particular devices or biological systems, which can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

Unless defined otherwise, all technical and scientific terms used herein have the same
15 meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. The following definitions supplement those in the art and are directed to the current application and are not to be imputed to any related or unrelated case, e.g., to any commonly owned patent or application. Although any methods and materials similar or equivalent to those described herein can be used in the practice for
20 testing of the invention, the preferred materials and methods are described herein. In describing and claiming the invention, the following terminology will be used in accordance with the definitions set out below.

As used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the content clearly dictates otherwise. Thus, for
25 example, reference to "a cell" includes a combination of two or more cells; reference to "bacteria" includes mixtures of bacteria, and the like.

As used herein, the terms "nucleic acid," "polynucleotide" and "oligonucleotide" refer to single-stranded or double-stranded nucleotide polymers comprised of more than two nucleotide subunits covalently joined together. The nucleotides may comprise
30 deoxyribonucleotides (containing 2-deoxy-D-ribose), ribonucleotides (containing D-ribose), and/or any other N-glycoside of a purine or pyrimidine base, or modified purine or pyrimidine bases, or any combination thereof. The sugar groups of the nucleotide subunits may also comprise modified derivatives of ribose or deoxyribose,

- 13 -

such as O-methyl ribose. The nucleotide subunits of an oligonucleotide may be joined by phosphodiester linkages, phosphorothioate linkages, methyl phosphonate linkages or by other linkages, including, but not limited to, rare or non-naturally-occurring linkages, that do not prevent hybridization of the oligonucleotide. Furthermore, an
5 oligonucleotide may have uncommon nucleotides or non-nucleotide moieties. With the addition of such analogs as locked nucleic acids (LNAs), the size of primers and probes can be reduced to as little as 8 nucleic acids. Locked nucleic acids are a novel class of bicyclic DNA analogs in which the 2' and 4' positions in the furanose ring are joined via an O-methylene (oxy-LNA), S-methylene (thio-LNA), or amino methylene (amino-
10 LNA) moiety.

Oligonucleotide probes and amplification oligonucleotides of a defined sequence may be produced by techniques known to those of ordinary skill in the art, such as by chemical or biochemical synthesis, and by *in vitro* or *in vivo* expression from recombinant nucleic acid molecules, *e.g.*, bacterial or retroviral vectors. As used herein, an oligonucleotide
15 does not consist of wild-type chromosomal DNA or the *in vivo* transcription products thereof.

Oligonucleotides which are primer and/or probe sequences, as described below, may comprise DNA, RNA or nucleic acid analogs such as uncharged nucleic acid analogs including, but not limited to, peptide nucleic acids (PNAs), which are disclosed in
20 International Patent Application WO 92/20702, or morpholino analogs, which are described in U.S. Pat. Nos. 5,185,444, 5,034,506, and 5,142,047, all of which are herein incorporated by reference in their entireties. Such sequences can routinely be synthesized using a variety of techniques currently available. For example, a sequence of DNA can be synthesized using conventional nucleotide phosphoramidite chemistry and
25 the instruments available from Applied Biosystems, Inc. (Foster City, Calif.); DuPont (Wilmington, Del.); or Milligen (Bedford, Mass.). Similarly, and when desirable, the sequences can be labeled using methodologies well known in the art such as described in U.S. Pat. Nos. 5,464,746; 5,424,414; and 4,948,882, all of which are herein incorporated by reference in their entireties. Oligonucleotides (including, *e.g.*, labeled or modified
30 oligos) can also be ordered from a variety of commercial sources known to persons of skill. Essentially any nucleic acid can be custom ordered from any of a variety of commercial sources, for example, The Midland Certified Reagent Company (www.mcrc.com), The Great American Gene Company (www.genco.com), ExpressGen Inc. (www.expressgen.com), and QIAGEN (<http://oligos.qiagen.com>), among many
35 others.

- 14 -

A nucleic acid, nucleotide, polynucleotide or oligonucleotide can comprise the five biologically occurring bases (adenine, guanine, thymine, cytosine and uracil) and/or bases other than the five biologically occurring bases. These bases may serve a number of purposes, e.g., to stabilize or destabilize hybridization; to promote or inhibit probe degradation; or as attachment points for detectable moieties or quencher moieties. For example, a polynucleotide of the invention can contain one or more modified, non-standard, or derivatized base moieties, including, but not limited to, N⁶-methyl-adenine, N⁶-tert-butyl-benzyl-adenine, imidazole, substituted imidazoles, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N⁶-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N⁶-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N⁶-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acidmethylene ester, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, 2,6-diaminopurine, and 5-propynyl pyrimidine. Other examples of modified, non-standard, or derivatized base moieties may be found in U.S. Patent Nos. 6,001,611, 5,955,589, 5,844,106, 5,789,562, 5,750,343, 5,728,525, and 5,679,785, each of which is incorporated herein by reference in its entirety.

Furthermore, a nucleic acid, nucleotide, polynucleotide or oligonucleotide can comprise one or more modified sugar moieties including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose. A nucleic acid, nucleotide, polynucleotide or oligonucleotide can comprise phosphodiester linkages or modified linkages including, but not limited to phosphotriester, phosphoramidate, siloxane, carbonate, carboxymethylene ester, acetamidate, carbamate, thioether, bridged phosphoramidate, bridged methylene phosphonate, phosphorothioate, methylphosphonate, phosphorodithioate, bridged phosphorothioate or sulfone linkages, and combinations of such linkages.

An "isolated" nucleic acid molecule, as used herein, is one that is separated from nucleotide sequences which normally flank the nucleic acid molecule and/or has been

completely or partially purified from other biological material (e.g., protein) normally associated with the nucleic acid.

- The nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated. Thus, recombinant DNA contained in a vector is included in the definition of "isolated" as used herein. Also, isolated nucleic acid molecules include recombinant DNA molecules in heterologous host cells, as well as partially or substantially purified DNA molecules in solution. "Isolated" nucleic acid molecules also encompass *in vivo* and *in vitro* RNA transcripts of the DNA molecules of the invention. An isolated nucleic acid molecule or nucleotide sequence can include a nucleic acid molecule or nucleotide sequence that is synthesized chemically or by recombinant means. Also, isolated polynucleotides include recombinant DNA molecules in heterologous organisms, as well as partially or substantially purified DNA molecules in solution. *In vivo* and *in vitro* RNA transcripts of the DNA molecules of the invention are also encompassed by "isolated" nucleotide sequences. Such polynucleotides are useful, e.g., as primers and/or probes for detecting polymorphisms, in the manufacture of the encoded polypeptide, as probes for isolating homologous sequences (e.g., from other mammalian species), for gene mapping (e.g., by *in situ* hybridization with chromosomes), or for detecting expression of the gene in tissue (e.g., human tissue), such as by Northern blot analysis.
- 20 The isolated nucleic acid molecules may be RNA, mRNA, DNA, or cDNA, for example, and may be double- or single-stranded. They may encode the sense strand, the non-coding regions, and/or the antisense strand. The nucleic acid molecule can include all or a portion of the coding sequence of the gene and can further comprise additional non-coding regions such as introns and non-coding 3' and 5' sequences (including regulatory sequences, for example). Additionally, the nucleic acid molecule can be fused to a marker sequence, for example, a sequence that can be used to purify the nucleic acid molecule.

- The nucleic acid molecules of the invention can comprise one or more modified nucleotide residues. The modification may be at the base, sugar and/or phosphate moiety and include, for example, halogenation, hydroxylation, alkylation, an attached linker and/or label. The modifications can further comprise, for example, labeling, methylation, internucleotide modifications such as uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates), charged linkages (e.g., phosphorothioates, phosphorodithioates), pendent moieties (e.g., polypeptides),
- 30

intercalators (e.g., acridine, psoralen), chelators, alkylators, and modified linkages (e.g., alpha anomeric nucleic acids). Also included are synthetic molecules that mimic nucleic acid molecules in the ability to bind to a designated sequence via hydrogen bonding and other chemical interactions. Such molecules include, for example, those in which
5 peptide linkages substitute for phosphate linkages in the backbone of the molecule.

In certain embodiments, nucleic acid molecules of the invention include, but are not limited to, FRZB mRNA, cDNA and/or genomic DNA molecules. Nucleic acid molecules of the invention also include oligonucleotides, e.g., an oligonucleotide comprising one or more of the FRZB SNPs described herein.

10 As used herein, the term "primer" refers to an oligonucleotide having a hybridization specificity sufficient for the initiation of an enzymatic polymerization under predetermined conditions, for example in an amplification technique such as polymerase chain reaction (PCR), in a process of sequencing, in a method of reverse transcription and/or the like.

15 As used herein, the term "probe" refers to an oligonucleotide having a hybridization specificity sufficient for binding to a defined target sequence under predetermined conditions, for example in an amplification technique such as a 5'-nuclease reaction, in a hybridization-dependent detection method, such as a Southern or Northern blot, and/or the like. Primers and probes may be used in a variety of ways and may be defined by the
20 specific use. For example, a "capture probe" is immobilized or can be immobilized on a solid support by any appropriate means, including, but not limited to: by covalent bonding, by adsorption, by hydrophobic and/or electrostatic interaction, or by direct synthesis on a solid support (see in particular patent application WO 92 10092). A "detection probe" may be labeled by means of a marker chosen, for example, from
25 radioactive isotopes, enzymes, in particular enzymes capable of acting on a chromogenic, fluorescent or luminescent substrate (in particular a peroxidase or an alkaline phosphatase), chromophoric chemical compounds, chromogenic, fluorogenic or luminescent compounds, analogues of nucleotide bases, and ligands such as biotin. Illustrative fluorescent compounds include, for example, fluorescein,
30 carboxyfluorescein, tetrachlorofluorescein, hexachlorofluorescein, Cy3, tetramethylrhodamine, Cy3.5, carboxy-x-rhodamine, Texas Red, Cy5, and Cy5.5. Illustrative luminescent compounds include, for example, luciferin and 2,3-dihydrophthalazinediones, such as luminol.

All of the oligonucleotides, primers and probes of the invention, whether hybridization assay probes, amplification primers, or other oligonucleotides, may be modified with chemical groups to enhance their performance or to facilitate the characterization of amplification products. For example, backbone-modified oligonucleotides such as those
5 having phosphorothioate or methylphosphonate groups which render the oligonucleotides resistant to the nucleolytic activity of certain polymerases or to nuclease enzymes may allow the use of such enzymes in an amplification or other reaction.

Another example of modification involves using non-nucleotide linkers (*e.g.*, Arnold, *et al.*, "Non- Nucleotide Linking Reagents for Nucleotide Probes", EP 0 313 219 hereby
10 incorporated by reference herein in its entirety) incorporated between nucleotides in the nucleic acid chain which do not interfere with hybridization or the elongation of the primer. Amplification oligonucleotides may also contain mixtures of the desired modified and natural nucleotides.

The 3' end of an amplification primer or a probe may optionally be blocked to prevent
15 initiation of DNA synthesis as described by McDonough, *et al.*, entitled "Nucleic Acid Sequence Amplification", WO94/03472 which enjoys common ownership with the invention and is hereby incorporated by reference herein in its entirety. A mixture of different 3' blocked amplification oligonucleotides, or of 3' blocked and unblocked oligonucleotides, may increase the efficiency of nucleic acid amplification, as described
20 therein. The 5' end of the oligonucleotides may be modified to be resistant to the 5'-exonuclease activity present in some nucleic acid polymerases. Such modifications can be carried out by adding a non-nucleotide group to the terminal 5' nucleotide of the primer using techniques such as those described by Arnold, *et al.*, *supra*, entitled "Non-Nucleotide Linking Reagents for Nucleotide Probes", incorporated by reference herein.

25 Once synthesized, selected oligonucleotide probes may be labeled by any of several well-known methods (*e.g.*, J. Sambrook, *infra*). Useful labels include radioisotopes as well as non-radioactive reporting groups. Isotopic labels include ^3H , ^{35}S , ^{32}P , ^{125}I , ^{57}Co and ^{14}C . Isotopic labels can be introduced into the oligonucleotide by techniques known in the art such as nick translation, end labeling, second strand synthesis, the use of reverse
30 transcription, and by chemical methods. When using radiolabeled probes hybridization can be detected, *e.g.*, by autoradiography, scintillation counting, or gamma counting. The detection method selected will depend upon the particular radioisotope used for labeling.

Non-isotopic materials can also be used for labeling and may be introduced internally into the nucleic acid sequence or at the end of the nucleic acid sequence. Modified nucleotides may be incorporated enzymatically or chemically. Chemical modifications of the probe may be performed during or after synthesis of the probe, for example, through the use of non-nucleotide linker groups as described by Arnold, *et al.*, *supra* "Non-Nucleotide Linking Reagents for Nucleotide Probes," incorporated by reference herein. Non-isotopic labels include fluorescent molecules; chemiluminescent molecules, enzymes, cofactors, enzyme substrates, haptens or other ligands.

In one embodiment, the probes are labeled with an acridinium ester. Acridinium ester labeling may be performed as described by Arnold et al., U.S. Pat. No. 5,185,439, entitled "Acridinium Ester Labeling and Purification of Nucleotide Probes," issued Feb. 9, 1993 and hereby incorporated by reference herein in its entirety.

The term "sequence-specific oligonucleotide" refers to an oligonucleotide that hybridizes (under defined conditions) to a target nucleic acid with a signal to noise ratio at least 2x higher than a signal to noise ratio at which it hybridizes to a nucleic acid that is not the target. For example, a sequence-specific oligonucleotide can hybridize, under stringent conditions, to a region of FRZB comprising a first allele of a SNP (the target nucleic acid) with a signal to noise ratio at least 2x (e.g., at least 5x or 10x) higher than a signal to noise ratio at which the oligonucleotide hybridizes to the region of FRZB comprising a second allele of the SNP.

The term "FRZB gene" or "FRZB locus" refers to the genomic nucleic acid sequence that encodes the FRZB protein. The nucleotide sequence of a gene, as used herein, encompasses coding regions, referred to as exons, intervening, non-coding regions, referred to as introns, and upstream and/or downstream regions. Upstream or downstream regions can include regions of the gene that are transcribed but not part of an intron or exon, or regions of the gene that comprise, for example, binding sites for factors that modulate gene transcription. The sequence for the human FRZB genomic sequence is provided at GenBank accession numbers NT_005100.3, NT_005265, and NT_005403, and a portion of the sequence is provided herein as SEQ ID NO:1. The sequence for the human FRZB mRNA is provided at GenBank accession number NM_001463, and is provided herein as SEQ ID NO:2.

The term "allele", as used herein, refers to a sequence variant of a gene and/or a polymorphism. Alleles are typically identified with respect to one or more polymorphic

positions, with the rest of the gene sequence unspecified. For example, a FRZB allele may be defined by the nucleotide present at a single SNP, or by the nucleotides present at a plurality of SNPs. Examples of such FRZB SNPs are provided in Table 1, below.

For convenience, the allele present at the higher or highest frequency in the population will be referred to as the wild-type allele, and less frequent allele(s) will be referred to as mutant allele(s). (However, it is worth noting that an allele which is more frequent in one population may be less frequent in a different population.) This designation of an allele as a mutant is meant solely to distinguish the allele from the wild-type allele and is not meant to indicate a change or loss of function.

10 The terms "polymorphic" and "polymorphism", as used herein, refer to the condition in which two or more variants of a specific genomic sequence, or the encoded amino acid sequence, can be found in a population. The terms refer either to the nucleic acid sequence or the encoded amino acid sequence; the use will be clear from the context. The term "polymorphic region" or "polymorphic site" refers to a region of the nucleic acid where the nucleotide difference that distinguishes the variants occurs, or, for amino acid sequences, a region of the amino acid sequence where the amino acid difference that distinguishes the protein variants occurs. As used herein, the term "single nucleotide polymorphism", or SNP, refers to polymorphism at a polymorphic site consisting of a single nucleotide position. As will be clear from the context, the term 20 polymorphism can refer to a specific variant sequence at a polymorphic site; for example, the term SNP can refer to the specific nucleotide (e.g., A, C, G, or T) occupying a polymorphic site consisting of a single nucleotide position.

Individual amino acids in a sequence are represented herein as AN or NA, wherein A is the amino acid in the sequence and N is the position in the sequence. In the case that 25 position N is polymorphic, it is convenient to designate one variant (e.g., the more frequent variant) as A₁N and the other variant (e.g., the less frequent variant) as NA₂. Alternatively, the polymorphic site, N, is represented as A₁NA₂, wherein A₁ is the amino acid in one variant and A₂ is the amino acid in the other variant. It is worth noting that an allele which is more frequent in one population may be less frequent in a different 30 population. Either the one-letter or three-letter codes are used for designating amino acids (see Lehninger, *BioChemistry 2nd ed.*, 1975, Worth Publishers, Inc. New York, NY: pages 73-75, incorporated herein by reference). The amino acid positions are numbered based on the sequence of the mature FRZB protein.

Representations of nucleotides and single nucleotide variations in DNA sequences are analogous to the representations of amino acids. For example, C18679T represents a single nucleotide polymorphism at nucleotide position 18679, wherein cytosine is present in the more frequent (wild-type) allele in the population and thymidine is present in the less frequent (mutant) allele. In general, a SNP can be represented as A_1NA_2 , wherein A_1 is the nucleotide present in one variant and A_2 is the nucleotide in the other variant. The single letter codes for the nucleotides are well known to those in the art, i.e., C for cytosine; A for adenine; T for thymidine, G for guanine, I for inosine, and U for uracil. It will be clear that in a double stranded form, the complementary strand of each allele will contain the complementary base at the polymorphic position; a SNP (or other polymorphism) can thus be described and/or detected with reference to the nucleotide(s) occupying the polymorphic site on either strand.

As used herein, the term "predisposing polymorphism" refers to a polymorphism that is positively associated with a condition, such as, for example, obesity and/or osteoporosis. The presence of a predisposing polymorphism in an individual could be indicative that the individual has an increased risk for the disease relative to an individual without the polymorphism. The term "protective polymorphism" refers to a polymorphism that is negatively associated with a condition. The presence of a protective polymorphism in an individual could be indicative that the individual has a decreased risk for the disease relative to an individual without the polymorphism.

The term "obesity-associated polymorphism" or "obesity-related polymorphism" refers to a polymorphism that is associated with obesity (e.g., with a high body mass index), either positively or negatively.

The term "osteoporosis-associated polymorphism" or "osteoporosis-related polymorphism" refers to a polymorphism that is associated with osteoporosis (e.g., with increased incidence of hip and/or vertebral fracture and/or decreased bone mineral density), either positively or negatively.

The term "association" or "associated with" in the context of this invention refers to the presence of a disease or phenotypic trait in individuals with one or more specific alleles or polymorphisms in one or more specific genes.

As used herein, the term "odds ratio" (OR) refers to the ratio of the frequency of the disease in individuals having a particular marker (allele or polymorphism) to the frequency of the disease in individuals without the marker (allele or polymorphism).

As used herein, the term "linkage disequilibrium" (LD) refers to alleles at different loci that are not associated at random, i.e., not associated in proportion to their frequencies. If the alleles are in positive linkage disequilibrium, then the alleles occur together more often than expected, assuming statistical independence. Conversely, if the alleles are in negative linkage disequilibrium, then the alleles occur together less often than expected assuming statistical independence.

The term "genotype", as used herein, refers to a description of the allele(s) of a gene or genes contained in an individual or a sample. As used herein, no distinction is made between the genotype of an individual and the genotype of a sample originating from the individual. Although, typically, a genotype is determined from samples of diploid cells, a genotype can be determined from a sample of haploid cells, such as a sperm cell. Similarly, an individual's "genotype at one or more polymorphic sites" refers to a description of the allele(s) of one or more polymorphisms contained in the individual or a sample. For example, an individual's genotype for a SNP is defined by the nucleotide present at that polymorphic site.

The term "haplotype", as used herein, refers to a description of the variants of a gene or genes contained on a single chromosome, i.e., the genotype of a single chromosome. A haplotype is a set of maternally inherited alleles, or a set of paternally inherited alleles, at any locus. A haplotype may also refer to two or more SNPs grouped together.

As used herein, the term "target region" refers to a region of a nucleic acid which is to be analyzed and usually includes at least one polymorphic region.

The term "stringent" as used herein refers to hybridization and wash conditions that are at or near the T_m for a particular sequence, taking into account considerations such as salt concentration and oligonucleotide length and base composition, for example.

Generally, stringent conditions are selected to be about 5 °C to 15 °C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Higher stringency conditions are optionally selected, for example, equal to the T_m or even 5 °C or more greater than the T_m . The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is at least about 0.02 molar at pH 7 and the temperature is at least about 50 °C for a sequence with a T_m of about 55-65 °C. As other factors may significantly affect the stringency of hybridization, including, among others, base composition, length of the nucleic acid

- 22 -

strands, the presence of organic solvents, and the extent of base mismatching, the combination of parameters is more important than the absolute measure of any one.

5 A "label" is a moiety that facilitates detection of a molecule. Common labels in the context of the present invention include fluorescent, luminescent, and/or colorimetric labels. Suitable labels include radionucleotides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents teaching the use of such labels include U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Many labels are commercially available and can be used in the context of the invention.

10 A "polynucleotide sequence", "nucleotide sequence", or "nucleic acid sequence" is a polymer of nucleotides (an oligonucleotide, a DNA, a nucleic acid, etc.) or a character string representing a nucleotide polymer, depending on context. From any specified polynucleotide sequence, either the given nucleic acid or the complementary polynucleotide sequence (e.g., the complementary nucleic acid) can be determined.

15 A variety of additional terms are defined or otherwise characterized herein.

DETAILED DESCRIPTION

20 Certain embodiments of the invention stem from the observation that at least one polymorphism in the FRZB gene is correlated with an individual's risk for obesity and/or osteoporosis. Further embodiments provide methods for detecting an individual's increased or decreased risk for obesity and/or osteoporosis. Further embodiments provide kits, reagents and arrays useful for detecting an individual's risk for obesity and/or osteoporosis.

25 In one aspect, the invention provides a method for detecting an individual's increased or decreased risk for obesity by detecting the presence of at least one obesity-associated polymorphism in the FRZB locus in a nucleic acid sample of the individual, wherein the presence of said at least one polymorphism indicates the individual's increased or decreased risk for obesity.

In a further aspect, the invention provides a method for detecting an individual's increased or decreased risk for osteoporosis by detecting the presence of at least one osteoporosis-associated polymorphism in the FRZB locus in a nucleic acid sample of the individual, wherein the presence of said at least one polymorphism indicates the individual's increased or decreased risk for osteoporosis.

In a further embodiment, the invention provides a method for detecting the presence of osteoporosis in an individual by a combination of the diagnostic test for predisposition and at least one other clinical test for osteoporosis, including but not limited to: a bone-turnover assay and any type of bone scan. Other diseases, conditions or criteria which may predispose for or be used in addition to the above include, but are not limited to, hyperthyroidism, posttransplantation, malabsorption, hyperparathyroidism, alcoholism, and family history.

In one embodiment, the FRZB polymorphism is selected from the FRZB polymorphisms listed in Table 1. In a further embodiment, more than one polymorphism is detected in the nucleic acid sample, at least one of which is selected from the polymorphisms listed in Table 1. In a further embodiment, at least two of the polymorphisms selected from those listed in Table 1 are detected. Examples of polymorphisms in FRZB that can be used include but are not limited to: T2303723C, C18679T, G19524A, T19575G, T22242A, G23043A, G23415A, T23549C, A24791G, C26794G, and G27014A.

The individual can be any mammal, including human, and belong to any race or population. The individual may be male or female. However, it is understood that the methods, kits and compositions described herein would be ideally suited for the analysis of osteoporosis in women over age 50, and/or post-menopausal women. Further, the methods, kits and compositions described herein would be ideally suited to the analysis of osteoporosis in men or women between the ages of 35 and 77. More particularly, men and women over 40 years of age would be ideally suited for analysis.

In a further embodiment, men and women between birth and the age of about 75 may be screened for a predisposition to obesity, more specifically, men and women between the ages of 5 and 55. Of interest would be to identify an individual predisposed to obesity at an early age and/or prenatally, in order to prevent the weight gain before it happens. In a further embodiment, the individual is identified at certain high-risk times or situations in the lifetime. Such times include but are not limited to: adolescence,

- 24 -

menopause, old age, periods of depression, illnesses which predispose to obesity and/or radical life changes. In addition, before starting on a diet regimen, it may be of interest to determine a patient's predisposition to obesity, which may influence the type of program selected, speed of weight loss, and the ability of the patient to maintain a different weight level.

The nucleic acid sample can be obtained from any part of the individual's body, including, but not limited to: hair, skin, nails, tissues, such as organs or tumors, or bodily fluids, such as saliva, blood, plasma, serum, spinal fluid, lymph, synovial fluid, semen, seminal fluid, bronchio-alveolar lavage, urine, or tears, as well as samples of isolated blood or tissue cells or in vitro cell culture constituents (including, but not limited to, conditioned medium obtained from the growth of cells in cell culture medium, recombinant cells and cell components). The nucleic acid sample can, but need not, be amplified by any amplification method including, but not limited to, polymerase chain reaction ("PCR").

The polymorphism can be any predisposing or protective polymorphism in the FRZB locus. In one embodiment of the invention, the polymorphism can be any polymorphism identified as predisposing or protective by methods taught herein. In one embodiment, the polymorphism can be a single nucleotide polymorphism (SNP) in the FRZB locus. In another embodiment, specific haplotypes in the FRZB locus as well as specific combinations of, and interactions between, SNPs at this locus can be indicative of an increased or a decreased risk of obesity and/or osteoporosis.

In a further embodiment, the presence of the polymorphism from only one parent is sufficiently predictive. In a further embodiment, the presence of the polymorphism from both parents is sufficiently predictive.

The polymorphism can be detected by any method known in the art for detecting the presence of a specific polymorphism in a nucleic acid sample. These methods include, but are not limited to, contacting the nucleic acid sample with one or more nucleic acid molecules that hybridize under stringent hybridization conditions to at least one FRZB polymorphism and detecting the hybridization, detection by amplification of the nucleic acid sample by, for example, PCR, and by direct sequencing of the nucleic acid sample.

In certain embodiments, an individual's risk for osteoporotic or obesity-related diseases is diagnosed from the individual's FRZB genotype. An individual who has at least one polymorphism statistically associated with osteoporosis and/or obesity possesses a factor

contributing to either an increased or a decreased risk as compared to an individual without the polymorphism. The statistical association of various FRZB polymorphisms (sequence variants) with obesity and/or osteoporosis is shown in the examples.

5 The genotype can be determined using any method capable of identifying nucleotide variation, e.g., nucleotide variation consisting of single nucleotide polymorphic sites. The particular method used is not a critical aspect of the invention. A number of suitable methods are described below.

10 In one embodiment of the invention, genotyping is carried out using oligonucleotide probes specific to variant FRZB sequences. In one embodiment, a region of the FRZB gene which encompasses one or several polymorphic sites of interest is amplified prior to, or concurrent with, the hybridization of probes directed to such sites. Probe-based assays for the detection of sequence variants are well known in the art.

15 Alternatively, genotyping is carried out using allele-specific amplification or extension reactions, wherein allele-specific primers are used which support primer extension only if the targeted allele is present. Typically, an allele-specific primer hybridizes to the FRZB gene such that the 3' terminal nucleotide aligns with a polymorphic position. Allele-specific amplification reactions and allele-specific extension reactions are well known in the art.

20 Another aspect of the invention relates to a kit useful for detecting the presence of a predisposing or a protective polymorphism in the FRZB locus in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed. The kit can comprise one or more oligonucleotides capable of detecting a predisposing or protective polymorphism in the FRZB locus as well as instructions for using the kit to detect susceptibility to obesity and/or osteoporosis. In preferred embodiments, the
25 oligonucleotide or oligonucleotides each individually comprise a sequence that hybridizes under stringent hybridization conditions to at least one FRZB polymorphism. In some embodiments, the oligonucleotide or oligonucleotides each individually comprise a sequence that is fully complementary to a nucleic acid sequence comprising a FRZB polymorphism.

30 In some embodiments, the oligonucleotide can be used to detect the presence of the FRZB polymorphism by hybridizing to the polymorphism under stringent hybridizing conditions. In some embodiments, the oligonucleotide can be used as an extension

primer in either an amplification reaction such as PCR or a sequencing reaction, wherein FRZB polymorphism is detected either by amplification or sequencing.

In certain embodiments, the kit can further comprise amplification or sequencing primers which can, but need not, be sequence-specific. The kit can also comprise reagents for labeling one or more of the oligonucleotides, or comprise labeled
5 oligonucleotides. Optionally, the kit can comprise reagents to detect the label.

In some embodiments, the kit can comprise one or more oligonucleotides that can be used to detect the presence of two or more predisposing or protective FRZB polymorphisms or combinations of predisposing polymorphisms, protective
10 polymorphisms or both.

In another aspect, the invention provides an array useful for detecting the presence of a predisposing or a protective FRZB polymorphism in a nucleic acid sample of an individual whose risk for obesity and/or osteoporosis is being assessed. The array can comprise one or more oligonucleotides capable of detecting a predisposing or protective
15 FRZB polymorphism. The oligonucleotides can be immobilized on a substrate, *e.g.*, a membrane or glass. In preferred embodiments, the oligonucleotide or oligonucleotides each individually comprise a sequence that can hybridize under stringent hybridization conditions to a nucleic acid sequence comprising a FRZB polymorphism. In some
20 embodiments, the oligonucleotide or oligonucleotides each individually comprise a sequence that is fully complementary to a nucleic acid sequence comprising a FRZB polymorphism. The oligonucleotide or oligonucleotides can, but need not, be labeled. In some embodiments, the array can be a microarray.

In some embodiments, the array can comprise one or more oligonucleotides used to detect the presence of two or more predisposing or protective FRZB polymorphisms or
25 combinations of predisposing polymorphisms, protective polymorphisms or both.

One aspect of the invention provides nucleic acids, for example, nucleic acids comprising one or more novel polymorphisms in the FRZB gene and/or nucleic acids useful for detecting one or more FRZB polymorphisms. Accordingly, one embodiment of the invention is an isolated nucleic acid molecule comprising a portion of the FRZB
30 gene, its complement, and/or a variant thereof. Preferably said variant comprises at least one of the polymorphisms identified herein. Even more preferably, said variant comprises at least one of the polymorphisms identified herein to be associated with obesity. Alternatively, said variant comprises at least one of the polymorphisms

identified herein to be associated with osteoporosis. Thus, in one embodiment, the nucleic acid molecule comprises at least one of the FRZB polymorphisms provided in Table 1. In a further embodiment, the nucleic acid molecule comprises or consists of a primer and/or a probe specific to at least one of the polymorphisms identified in the FRZB gene (e.g., those identified herein to be associated with obesity and/or

5 osteoporosis).

SNPS

In one aspect, the invention provides a method for detecting an individual's increased or decreased risk for obesity by detecting the presence of one or more FRZB SNPs in a

10 nucleic acid sample of the individual, wherein the presence of said SNP(s) indicates the individual's increased or decreased risk for obesity. In a further aspect, the invention provides a method for detecting an individual's increased or decreased risk for osteoporosis by detecting the presence of one or more FRZB SNPs in a nucleic acid

15 sample of the individual, wherein the presence of said SNP(s) indicates the individual's increased or decreased risk for osteoporosis. The SNPs can be any SNPs in the FRZB locus including SNPs in exons, introns and/or upstream and/or downstream regions. Examples of such SNPs include, but are not limited to, those provided in Table 1, below, and discussed in detail in the Examples. In one embodiment, the SNPs present in the

20 FRZB locus are identified by genotyping the FRZB SNPs.

Table 1: FRZB SNPs

FRZB SNP	Standard Name	Position in NT_005403	SNP Source	Pos. in Gene/Change
FRZB1_T2303723C	NT_005265.11_2303723	NT_005403_33933138	rs6433992	Intron 1/T-C
FRZB_C18679T	NT_005100.3_18679	NT005403_33917116	rs288330	Intron 2/C-T
FRZB_G19524A	NT_005100.3_19524	NT005403_33916271	rs2242070	Intron 3/G-A
FRZB_T19575G	NT_005100.3_19575	NT005403_33916220	RMS SNP	Intron 3/T-G
FRZB_T22242A	NT_005100.3_22242	NT005403_33913553	rs288327	Intron 3/T-A
FRZB_G23043A	NT_005100.3_23043	NT005403_33912752	rs288326	Exon 4/G-A->Arg200Trp
FRZB_G23415A	NT_005100.3_23415	NT005403_33912380	rs1561369	Intron 4/G-A
FRZB_T23549C	NT_005100.3_23549	NT005403_33912246	rs288325	Intron 4/T-C

FRZB SNP	Standard Name	Position in NT_005403	SNP Source	Pos. in Gene/Change
FRZB_A24791G	NT_005100.3_24791	NT005403_33911004	rs288324	Intron 5/A-G
FRZB_C26794G	NT_005100.3_26794	NT005403_33909000	rs7775	Exon 6/C-G - >Arg324Gly
FRZB_G27014A	NT_005100.3_27014	NT005403_33908780	rs13009	3' UTR/G-A

In certain embodiments, the genotype of one FRZB SNP can be used to determine an individual's risk for obesity and/or osteoporotic disease. In other embodiments, the genotypes of a plurality of FRZB SNPs can be used. In other embodiments, certain combinations of SNPs at either the same or different loci can be used. It is to be understood that some of the SNPs related to an increased or decreased risk of obesity may be the same as those related to either an increased or decreased risk of osteoporosis.

GENOTYPING METHODS

- 10 In the methods of the invention, the alleles present in a sample are identified by identifying the nucleotide present at one or more of the polymorphic sites in a nucleic acid sample of an individual. A number of methods are known in the art for identifying the nucleotide present at polymorphic sites. The particular method used to identify the genotype is not a critical aspect of the invention. Although considerations of
- 15 performance, cost, and convenience will make particular methods more desirable than others, it will be clear that any method that can identify the nucleotide present will provide the information needed to identify the genotype. Preferred genotyping methods involve DNA sequencing, allele-specific amplification, or probe-based detection of amplified nucleic acid.
- 20 FRZB alleles can be identified by DNA sequencing methods, such as the chain termination method (Sanger *et al.*, 1977, *Proc. Natl. Acad. Sci.*, 74:5463-5467, incorporated herein by reference), which are well known in the art. In one embodiment, a portion of the gene encompassing the polymorphic site is amplified and either cloned into a suitable plasmid and then sequenced, or sequenced directly. PCR-
- 25 based sequencing is described in U.S. Patent No. 5,075,216; Brow, in *PCR Protocols*, 1990, (Innis *et al.*, eds., Academic Press, San Diego), chapter 24; and Gyllensten, in *PCR Technology*, 1989 (Erich, ed., Stockton Press, New York), chapter 5; each incorporated herein by reference. Typically, sequencing is performed using an automated DNA

sequencer, which are commercially available from, for example, PE Biosystems (Foster City, CA), Pharmacia (Piscataway, NJ), Genomyx Corp. (Foster City, CA), LI-COR Biotech (Lincoln, NE), GeneSys technologies (Sauk City, WI), and Visible Genetics, Inc. (Toronto, Canada).

- 5 FRZB alleles can also be identified using amplification-based genotyping methods. Various nucleic acid amplification methods known in the art can be used in to detect nucleotide changes in a target nucleic acid. A preferred method is the polymerase chain reaction (PCR), which is now well known in the art, and described in U.S. Patent Nos. 4,683,195; 4,683,202; and 4,965,188; each incorporated herein by reference. Examples of
10 the numerous articles published describing methods and applications of PCR are found in PCR Applications, 1999, (Innis *et al.*, eds., Academic Press, San Diego), PCR Strategies, 1995, (Innis *et al.*, eds., Academic Press, San Diego); PCR Protocols, 1990, (Innis *et al.*, eds., Academic Press, San Diego); and PCR Technology, 1989, (Erich, ed., Stockton Press, New York); each incorporated herein by reference. Commercial
15 vendors, such as PE Biosystems (Foster City, CA) market PCR reagents and publish PCR protocols.

- Other suitable amplification methods include the ligase chain reaction (Wu and Wallace, 1988, *Genomics* 4:560-569); the strand displacement assay (Walker *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:392-396, Walker *et al.* 1992, *Nucleic Acids Res.* 20:1691-
20 1696, and U.S. Patent No. 5,455,166); and several transcription-based amplification systems, including the methods described in U.S. Patent Nos. 5,437,990; 5,409,818; and 5,399,491; the transcription amplification system (TAS) (Kwoh *et al.*, 1989, *Proc. Natl. Acad. Sci. USA*, 86:1173-1177); and self-sustained sequence replication (3SR) (Guatelli *et al.*, 1990, *Proc. Natl. Acad. Sci. USA*, 87:1874-1878 and WO 92/08800); each
25 incorporated herein by reference. Alternatively, methods that amplify the probe to detectable levels can be used, such as Q β -replicase amplification (Kramer *et al.*, 1989, *Nature*, 339:401-402, and Lomeli *et al.*, 1989, *Clin. Chem.*, 35:1826-1831, both of which are incorporated herein by reference). A review of known amplification methods is provided in Abramson *et al.*, 1993, *Current Opinion in Biotechnology*, 4:41-47,
30 incorporated herein by reference.

Genotyping can also be carried out by detecting and analyzing FRZB mRNA under conditions when both maternal and paternal chromosomes are transcribed. Amplification of RNA can be carried out by first reverse-transcribing the target RNA using, for example, a viral reverse transcriptase, and then amplifying the resulting

cDNA, or using a combined high-temperature reverse-transcription-polymerase chain reaction (RT-PCR), as described in U.S. Patent Nos. 5,310,652; 5,322,770; 5,561,058; 5,641,864; and 5,693,517; each incorporated herein by reference (see also Myers and Sigua, 1995, in PCR Strategies, *supra*, chapter 5).

- 5 FRZB alleles can also be identified using allele-specific amplification or primer extension methods, which are based on the inhibitory effect of a terminal primer mismatch on the ability of a DNA polymerase to extend the primer. To detect an allele sequence using an allele-specific amplification or extension-based method, a primer complementary to the FRZB genes is chosen such that the 3' terminal nucleotide hybridizes at the polymorphic
10 position. In the presence of the allele to be identified, the primer matches the target sequence at the 3' terminus and primer is extended. In the presence of only the other allele, the primer has a 3' mismatch relative to the target sequence and primer extension is either eliminated or significantly reduced. Allele-specific amplification- or extension-based methods are described in, for example, U.S. Patent Nos. 5,137,806; 5,595,890;
15 5,639,611; and U.S. Patent No. 4,851,331, each incorporated herein by reference.

Using allele-specific amplification-based genotyping, identification of the alleles requires only detection of the presence or absence of amplified target sequences. Methods for the detection of amplified target sequences are well known in the art. For example, gel electrophoresis (see Sambrook *et al.*, 1989, *infra*) and the probe hybridization assays
20 described above have been used widely to detect the presence of nucleic acids.

Allele-specific amplification-based methods of genotyping can facilitate the identification of haplotypes, as described in the examples. Essentially, the allele-specific amplification is used to amplify a region encompassing multiple polymorphic sites from only one of the two alleles in a heterozygous sample. The SNP variants present within
25 the amplified sequence are then identified, such as by probe hybridization or sequencing.

An alternative probe-less method, referred to herein as a kinetic-PCR method, in which the generation of amplified nucleic acid is detected by monitoring the increase in the total amount of double-stranded DNA in the reaction mixture, is described in Higuchi
30 *et al.*, 1992, *Bio/Technology*, 10:413-417; Higuchi *et al.*, 1993, *Bio/Technology*, 11:1026-1030; Higuchi and Watson, in PCR Applications, *supra*, Chapter 16; U.S. Patent Nos. 5,994,056 and 6,171,785; and European Patent Publication Nos. 487,218 and 512,334, each incorporated herein by reference. The detection of double-stranded target DNA

relies on the increased fluorescence that DNA-binding dyes, such as ethidium bromide or SYBR™ Green, exhibit when bound to double-stranded DNA. The increase of double-stranded DNA resulting from the synthesis of target sequences results in an increase in the amount of dye bound to double-stranded DNA and a concomitant detectable increase in fluorescence. For genotyping using the kinetic-PCR methods, amplification reactions are carried out using a pair of primers specific for one of the alleles, such that each amplification can indicate the presence of a particular allele. For example, by performing two amplifications, one using primers specific for the wild-type allele and one using primers specific for the mutant allele, the genotype of the sample with respect to that SNP can be determined. Similarly, by carrying out four amplifications, each with one of the possible pairs possible using allele specific primers for both the upstream and downstream primers, the genotype of the sample with respect to two SNPs can be determined. This gives haplotype information for a pair of SNPs.

Alleles can be also identified using probe-based methods, which rely on the difference in stability of hybridization duplexes formed between a probe and its corresponding target sequence comprising a FRZB allele. Under sufficiently stringent hybridization conditions, stable duplexes are formed only between a probe and its target allele sequence and not other allele sequences. The presence of stable hybridization duplexes can be detected by any of a number of well known methods. In general, it is preferable to amplify a nucleic acid encompassing a polymorphic site of interest prior to hybridization in order to facilitate detection. However, this is not necessary if sufficient nucleic acid can be obtained without amplification.

A probe suitable for use in the probe-based methods of the invention, which contains a hybridizing region either substantially complementary or exactly complementary to a target region of the FRZB gene or the complement thereof, wherein the target region encompasses the polymorphic site, and exactly complementary to one of the two allele sequences at the polymorphic site, can be selected using the guidance provided herein and well known in the art. Similarly, suitable hybridization conditions (e.g., stringent hybridization conditions), which depend on the exact size and sequence of the probe, can be selected empirically using the guidance provided herein and well known in the art (see, e.g., *Nucleic Acid Hybridization* (B.D. Hames and S.J. Higgins. eds., 1984) and Sambrook et al., *Molecular Cloning - A Laboratory Manual* (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 2000). The use of oligonucleotide probes to detect nucleotide variations including single base pair differences in sequence is described in, for example, Conner *et al.*, 1983, *Proc. Natl.*

Acad. Sci. USA, 80:278-282, and U.S. Patent Nos. 5,468,613 and 5,604,099, each incorporated herein by reference.

In some embodiments of the probe-based methods for determining the FRZB genotypes, multiple nucleic acid sequences from the FRZB genes which encompass the polymorphic sites are amplified and hybridized to a set of probes under sufficiently stringent hybridization conditions. The alleles present are inferred from the pattern of binding of the probes to the amplified target sequences. In this embodiment, amplification is carried out in order to provide sufficient nucleic acid for analysis by probe hybridization. Thus, primers are designed such that regions of the FRZB genes encompassing the polymorphic sites are amplified regardless of the allele present in the sample. Allele-independent amplification is achieved using primers which hybridize to conserved regions of the FRZB genes. The FRZB genes contain many invariant or monomorphic regions, and suitable allele-independent primers can be selected routinely from SEQ ID NO:1 or GenBank accession numbers NT_005100.3, NT_005265, or NT_005403.

Suitable assay formats for detecting hybrids formed between probes and target nucleic acid sequences in a sample are known in the art and include the immobilized target (dot-blot) format and immobilized probe (reverse dot-blot or line-blot) assay formats. Dot blot and reverse dot blot assay formats are described in U.S. Patent Nos. 5,310,893; 5,451,512; 5,468,613; and 5,604,099; each incorporated herein by reference.

In a dot-blot format, amplified target DNA is immobilized on a solid support, such as a nylon membrane. The membrane-target complex is incubated with labeled probe under suitable hybridization conditions, unhybridized probe is removed by washing under suitably stringent conditions, and the membrane is monitored for the presence of bound probe. A preferred dot-blot detection assay is described in the examples.

In the reverse dot-blot (or line-blot) format, the probes are immobilized on a solid support, such as a nylon membrane or a microtiter plate. The target DNA is labeled, typically during amplification by the incorporation of labeled primers. One or both of the primers can be labeled. The membrane-probe complex is incubated with the labeled amplified target DNA under suitable hybridization conditions, unhybridized target DNA is removed by washing under suitably stringent conditions, and the membrane is monitored for the presence of bound target DNA. A preferred reverse line-blot detection assay is described in the examples.

Probe-based genotyping can be carried out using a 5'-nuclease assay, as described in U.S. Patent Nos. 5,210,015; 5,487,972; and 5,804,375; and Holland *et al.*, 1988, *Proc. Natl. Acad. Sci. USA*, 88:7276-7280, each incorporated herein by reference. In the 5'-nuclease assay, labeled detection probes that hybridize within the amplified region are added during the amplification reaction mixture. The probes are modified so as to prevent the probes from acting as primers for DNA synthesis. The amplification is carried out using a DNA polymerase that possesses 5' to 3' exonuclease activity, *e.g.*, *Tth* DNA polymerase. During each synthesis step of the amplification, any probe which hybridizes to the target nucleic acid downstream from the primer being extended is degraded by the 5' to 3' exonuclease activity of the DNA polymerase. Thus, the synthesis of a new target strand also results in the degradation of a probe, and the accumulation of degradation product provides a measure of the synthesis of target sequences.

Any method suitable for detecting degradation product can be used in the 5'-nuclease assay. In a preferred method, the detection probes are labeled with two fluorescent dyes, one of which is capable of quenching the fluorescence of the other dye. The dyes are attached to the probe, preferably one attached to the 5' terminus and the other is attached to an internal site, such that quenching occurs when the probe is in an unhybridized state and such that cleavage of the probe by the 5' to 3' exonuclease activity of the DNA polymerase occurs in between the two dyes. Amplification results in cleavage of the probe between the dyes with a concomitant elimination of quenching and an increase in the fluorescence observable from the initially quenched dye. The accumulation of degradation product is monitored by measuring the increase in reaction fluorescence. U.S. Patent Nos. 5,491,063 and 5,571,673, both incorporated herein by reference, describe alternative methods for detecting the degradation of probe which occurs concomitant with amplification.

The 5'-nuclease assay can be used with allele-specific amplification primers such that the probe is used only to detect the presence of amplified product. Such an assay is carried out as described for the kinetic-PCR-based methods described above. Alternatively, the 5'-nuclease assay can be used with a target-specific probe.

Examples of other techniques that can be used for probe-based genotyping include, but are not limited to, Amplifluor™, Dye Binding-Intercalation, Fluorescence Resonance Energy Transfer (FRET), Hybridization Signal Amplification Method (HSAM), HYB Probes™, Invader/Cleavase Technology (Invader/CFLP™), Molecular Beacons™,

Origen™, DNA-Based Ramification Amplification (RAM™), Rolling circle amplification (RCA™), Scorpions™, Strand displacement amplification (SDA).

5 The assay formats described above typically utilize labeled oligonucleotides to facilitate detection of the hybrid duplexes. Oligonucleotides can be labeled by incorporating a label detectable by spectroscopic, photochemical, biochemical, immunochemical, radiological, radiochemical or chemical means. Useful labels include ³²P, fluorescent dyes, electron-dense reagents, enzymes (as commonly used in ELISAs), biotin, or haptens and proteins for which antisera or monoclonal antibodies are available. Labeled oligonucleotides of the invention can be synthesized and labeled using the techniques described above for synthesizing oligonucleotides. For example, a dot-blot assay can be carried out using probes labeled with biotin, as described in Levenson *et al.*, 1989, in *PCR Protocols: A Guide to Methods and Applications* (Innis *et al.*, eds., Academic Press. San Diego), pages 99-112, incorporated herein by reference. Following hybridization of the immobilized target DNA with the biotinylated probes under sequence-specific conditions, probes which remain bound are detected by first binding the biotin to 15 avidin-horseradish peroxidase (A-HRP) or streptavidin-horseradish peroxidase (SA-HRP), which is then detected by carrying out a reaction in which the HRP catalyzes a color change of a chromogen.

20 Whatever the method for determining which oligonucleotides of the invention selectively hybridize to FRZB allelic sequences in a sample, the central feature of the typing method involves the identification of the FRZB alleles present in the sample by detecting the variant sequences present. Further details on genotyping of SNPs are available in the literature; see, e.g., Lindblad-Toh *et al.*, 2000, *Nature Genetics* 24:381-386; Plant Genotyping: The DNA Fingerprinting of Plants, 2001, CABI Publishing; 25 Syvanen, 2001, *Nat. Rev. Genet.* 2:930-942; Kuklin *et al.*, 1998, *Genetic Testing* 1: 201-206; Gut, 2001, *Hum. Mutat.* 17:475-492; Ahmadian *et al.*, 2000, *Anal. Biochem.* 280:103-110; Useche *et al.*, 2001, *Genome Inform Ser Workshop Genome Inform* 12:194-203; Pastinen *et al.*, 2000, *Genome Res.* 10:1031-1042; Hacia, 1999, *Nature Genet.* 22:164-167; and Chen *et al.*, 2000, *Genome Res.* 10:549-557.

OTHER MARKERS

Other genetic markers and methods of detecting sequence polymorphisms are known in the art and can be applied to the practice of the present invention, including, but not limited to, restriction fragment length polymorphisms (RFLPs), random amplified polymorphic DNA (RAPD), arbitrary fragment length polymorphisms (AFLPs), simple sequence repeats (SSRs), single-stranded conformation polymorphisms (SSCPs), and amplified variable sequences. Discovery, detection, and genotyping of these and other types of genetic markers has been well described in the literature. See, e.g., Orita *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 86:2766-2770; USPN 6,399,855; Henry, ed., 2001, Plant Genotyping. The DNA Fingerprinting of Plants Wallingford: CABI Publishing; Phillips and Vasil, eds., 2001, DNA-based Markers in Plants Dordrecht: Kluwer Academic Publishers; Pejic *et al.*, 1998, *Theor. App. Genet.* 97:1248-1255; Bhatramakki *et al.*, 2002, *Plant Mol. Biol.* 48:539-47; Nickerson *et al.*, 1997, *Nucleic Acids Res.* 25:2745-2751; Underhill *et al.*, 1997, *Genome Res.* 7:996-1005; Shi, 2001, *Clin. Chem.* 47:164-172; Kwok, 2000, *Pharmacogenomics* 1:95-100; Rafalski *et al.*, 2002, *Cell Mol Biol Lett* 7:471-5; Ching and Rafalski, 2002, *Cell Mol Biol Lett.* 7:803-10; Powell *et al.*, 1996, *Mol. Breeding* 2:225-238; Vos *et al.*, 1995, *Nucl. Acids Res.* 23:4407; Becker *et al.*, 1995, *Mol. Gen. Genet.* 249:65; Meksem *et al.*, 1995, *Mol. Gen. Genet.* 249:74; Huys *et al.*, 1996, *Int'l J. Systematic Bacteriol.* 46:572; Jacob *et al.*, 1991, *Cell* 67:213; Taramino and Tingey, 1996, *Genome* 39:277-287; Condit and Hubbell, 1991, *Genome* 34:66; and Zietkiewicz *et al.*, 1994, *Genomics* 20:176-83.

ASSOCIATION ANALYSIS

Evaluation of the candidate gene FRZB for association with various phenotypes pertaining to obesity and osteoporosis is described in the Examples. In addition, design and execution of various types of association studies have been described in the art; see, e.g., Rao and Province, eds., 2001, *Advances in Genetics* volume 42, Genetic Dissection of Complex Traits; Balding *et al.*, eds., 2001, Handbook of Statistical Genetics, John Wiley and Sons Ltd.; Borecki and Suarez, 2001, *Adv Genet* 42:45-66; Cardon and Bell, 2001, *Nat Rev Genet* 2:91-99; and Risch, 2000, *Nature* 405:847-856. Association studies have been used both to evaluate candidate genes for association with a phenotypic trait (e.g., Thornsberry *et al.*, 2001, *Nature Genetics* 28:286-289) and to perform whole genome scans to identify genes that contribute to phenotypic variation.

KITS

The invention also relates to a kit comprising a container unit and components for practicing the present method. A kit can contain oligonucleotide probes specific for FRZB alleles as well as instructions for their use to determine risk for obesity and/or osteoporosis. In some cases, a kit may comprise detection probes fixed to an appropriate support membrane. The kit can also contain amplification primers for amplifying regions of the FRZB locus encompassing the polymorphic sites, as such primers are useful in the preferred embodiment of the invention. Alternatively, useful kits can contain a set of primers comprising an allele-specific primer for the specific amplification of FRZB alleles. Other optional components of the kits include additional reagents used in the genotyping methods as described herein. For example, a kit additionally can contain an agent to catalyze the synthesis of primer extension products, substrate nucleoside triphosphates, reagents for labeling and/or detecting nucleic acid (for example, an avidin-enzyme conjugate and enzyme substrate and chromogen if the label is biotin) and appropriate buffers for amplification or hybridization reactions.

One aspect of the invention provides kits for detecting presence of a first predisposing or protective polymorphism in an FRZB gene, e.g., in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed. Thus, one general class of embodiments provides a kit including one or more first oligonucleotides capable of detecting the first polymorphism and instructions for detecting the first polymorphism with the one or more first oligonucleotides and for correlating said detection to the individual's risk for osteoporosis and/or obesity, packaged in one or more containers.

Essentially all of the features noted for the method embodiments above apply to this embodiment as well, as relevant. For example, in a preferred class of embodiments, the first polymorphism is a single nucleotide polymorphism, e.g., a SNP selected from the group consisting of: the T allele of T2303723C, the C allele of T2303723C, the C allele of C18679T, the T allele of C18679T, the G allele of G19524A, the A allele of G19524A, the T allele of T22242A, the A allele of T22242A, the A allele of A24791G, the G allele of A24791G, the C allele of C26794G, the G allele of C26794G, the G allele of G27014A, and the A allele of G27014A. Other potential SNPs include, but are not limited to, either allele of T19575G, G23043A, G23415A, and T23549C.

In one aspect, the kit can be used to detect the presence of the first polymorphism by hybridization of a nucleic acid probe to the polymorphism. Thus, in one class of

- 37 -

embodiments, the one or more first oligonucleotides comprise at least one probe. In certain embodiments, the first oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the first polymorphism. In one class of embodiments, the first polymorphism is a first single nucleotide polymorphism comprising a first nucleotide at a first nucleotide position. In this class of embodiments, under stringent conditions, the first oligonucleotide hybridizes to a region of the FRZB gene comprising the first single nucleotide polymorphism with a signal to noise ratio that is at least 2x (e.g., at least 5x or at least 10x) the signal to noise ratio at which the first oligonucleotide hybridizes to the region of the FRZB gene comprising a second nucleotide at the first nucleotide position. The first oligonucleotide is typically fully complementary to the region of the FRZB gene comprising the first polymorphism, and typically comprises at least about 10 contiguous nucleotides complementary to the FRZB gene.

To facilitate detection of the polymorphism (e.g., through detection of hybridization between the one or more first oligonucleotides and a nucleic acid comprising the polymorphism), for example, the one or more first oligonucleotides optionally comprise a label, e.g., an isotopic, fluorescent, fluorogenic, luminescent or colorimetric label. In some embodiments, the label itself directly produces a detectable signal (e.g., a fluorescent label). In other embodiments, the kit also includes a reagent that detects the label (e.g., an enzyme that cleaves a colorimetric label, a binding moiety, or the like).

In one aspect, the one or more first oligonucleotides comprise one or more primers. The primer(s) can be used to detect the polymorphism, e.g., in an allele-specific amplification or extension reaction. For example, in one class of embodiments, the first polymorphism is a first single nucleotide polymorphism comprising a first nucleotide at a first nucleotide position, and the 3' nucleotide of one of the one or more first oligonucleotides is complementary to the first nucleotide.

The primer(s) can be used to amplify a region of FRZB comprising the polymorphism, e.g., for subsequent detection of the polymorphism by hybridization, sequencing, or the like. In one class of embodiments, the one or more first oligonucleotides comprise amplification primers, wherein the amplification primers amplify a nucleic acid sequence comprising the first polymorphism. In a related class of embodiments, the one or more first oligonucleotides comprise sequencing primers that flank the first polymorphism.

The one or more first oligonucleotides are optionally immobilized on a substrate. The substrate can be, for example, a planar substrate or a beaded substrate. The oligonucleotide(s) can be arranged in an array of other oligonucleotides used to detect other polymorphisms, e.g., other polymorphisms in FRZB.

- 5 The kit can optionally be used to detect more than one polymorphism (simultaneously or sequentially). Thus, in one class of embodiments, the kit also includes one or more second oligonucleotides capable of detecting a second polymorphism (and optionally third, fourth, fifth, etc. oligonucleotides capable of detecting third, fourth, fifth, etc. polymorphisms). The second polymorphism can be at the same polymorphic site as the
0 first or at a different polymorphic site (in FRZB or a different gene), and can be protective or predisposing.

ARRAYS AND SYSTEMS

- The invention also relates to an array, a support with immobilized oligonucleotides
5 useful for practicing the present method. A useful array can contain oligonucleotide probes specific for FRZB alleles or certain combinations of FRZB alleles. The oligonucleotides can be immobilized on a substrate, e.g., a membrane or glass. The oligonucleotides can, but need not, be labeled. In some embodiments, the array can be a micro-array. In some embodiments, the array can comprise one or more
10 oligonucleotides used to detect the presence of two or more FRZB alleles or certain combinations of FRZB alleles.

- One general class of embodiments provides arrays for detecting presence of one or more predisposing and/or protective polymorphisms in an FRZB gene, e.g., in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed. In
15 one class of embodiments, the array comprises a substrate and a plurality of oligonucleotides, each of which oligonucleotides hybridizes to a region of the FRZB gene comprising at least one of the polymorphisms. The hybridization detects the presence of the polymorphism, and this detection provides an indication of the individual's risk for osteoporosis and/or obesity. Typically, the array is used for detecting the presence of a
20 plurality of polymorphisms, e.g., multiple alleles at a single polymorphic site and/or different polymorphic sites.

Essentially all of the features noted for the method and kit embodiments above apply to this embodiment as well, as relevant. For example, the one or more polymorphisms

- 39 -

preferably comprise one or more single nucleotide polymorphisms. For example, at least one of the one or more polymorphisms can be selected from the group consisting of: the T allele of T2303723C, the C allele of T2303723C, the C allele of C18679T, the T allele of C18679T, the G allele of G19524A, the A allele of G19524A, the T allele of T22242A, the A allele of T22242A, the A allele of A24791G, the G allele of A24791G, the C allele of C26794G, the G allele of C26794G, the G allele of G27014A, and the A allele of G27014A. Other potential SNPs include, but are not limited to, either allele of T19575G, G23043A, G23415A, and T23549C.

In one class of embodiments in which the array can be used to detect presence of one or more SNPs, each of the oligonucleotides in the array hybridizes under stringent conditions to a region of the FRZB gene comprising one of the single nucleotide polymorphisms with a signal to noise ratio that is at least 2x (e.g., at least 5x or at least 10x) that at which the oligonucleotide hybridizes to a region of the FRZB gene comprising any of the remaining single nucleotide polymorphisms. Typically, one oligonucleotide is used to detect one SNP; that is, each of the oligonucleotides typically hybridizes to a distinct single nucleotide polymorphism.

As noted, the plurality of oligonucleotides are immobilized on a substrate, e.g., a planar substrate, a membrane, a glass slide, or the like. Typically, each of the plurality of oligonucleotides is immobilized at a known, pre-determined position on the substrate.

To facilitate detection of polymorphisms by specific hybridization with the oligonucleotides, each of the plurality of oligonucleotides is typically fully complementary to a region of the FRZB gene comprising one of the polymorphisms, and each of the plurality of oligonucleotides typically comprises at least about 10 contiguous nucleotides complementary to the FRZB gene. Each of the plurality of oligonucleotides optionally comprises a label, e.g., a label that facilitates detection of hybridization between the oligonucleotides and the polymorphisms.

The array is optionally part of a system. Thus, one class of embodiments provides a system comprising an array of the invention and system instructions that correlate the detection of the presence of one or more predisposing or protective polymorphisms to the individual's risk for osteoporosis and/or obesity. Systems, e.g., digital systems, are described in greater detail below.

In an array on a substrate, each oligonucleotide is typically bound (e.g., electrostatically or covalently bound, directly or via a linker) to the substrate at a unique location.

- 40 -

Methods of making, using, and analyzing such arrays (e.g., microarrays) are well known in the art. See, e.g., Wang *et al.*, 1998, *Science* 280:1077-82; Lockhart and Winzeler, 2000, *Nature* 405:827-836; and Scherf *et al.*, 2000, *Nat Genet.* 24:236-44. Arrays can be formed (e.g., printed), for example, using commercially available instruments such as a
5 GMS 417 Arrayer (Affymetrix, Santa Clara, CA). Suitable solid supports are commercially readily available. For example, a variety of membranes (e.g., nylon, PVDF, and nitrocellulose membranes) are commercially available, e.g., from Sigma-Aldrich, Inc. (www.sigmaaldrich.com). As another example, surface-modified and pre-coated slides with a variety of surface chemistries are commercially available, e.g., from
10 TeleChem International (www.arrayit.com), Corning, Inc. (Corning, NY), or Greiner Bio-One, Inc. (www.greinerbiooneinc.com). For example, silanated and silylated slides with free amino and aldehyde groups, respectively, are available and permit covalent coupling of molecules (e.g., oligos) to the slides. Slides with surface streptavidin are available and can bind biotinylated oligos. In addition, services that produce arrays of
15 nucleic acids of the customer's choice are commercially available, e.g., from TeleChem International (www.arrayit.com).

DIGITAL SYSTEMS

In general, various automated systems can be used to perform some or all of the method
20 steps as noted herein. In addition to practicing some or all of the method steps herein, digital or analog systems, e.g., comprising a digital or analog computer, can also control a variety of other functions such as a user viewable display (e.g., to permit viewing of method results by a user) and/or control of output features.

For example, certain of the methods described above are optionally implemented via a
25 computer program or programs (e.g., that correlate detection of the presence of one or more predisposing or protective polymorphisms to an individual's risk for osteoporosis and/or obesity). Thus, the present invention provides digital systems, e.g., computers, computer readable media, and/or integrated systems comprising instructions (e.g., embodied in appropriate software) for performing the methods herein. For example, a
30 digital system comprising instructions for correlating detection of the presence of one or more predisposing or protective polymorphisms to an individual's risk for osteoporosis and/or obesity, as described herein, is a feature of the invention. The digital system can also include information (data) corresponding to individual genotypes for a set of genetic markers, phenotypic values, and/or the like. The system can also aid in detection

of the one or more polymorphisms (e.g., by controlling a microarray scanner, or the like).

Standard desktop applications such as word processing software (e.g., Microsoft Word™ or Corel WordPerfect™) and/or database software (e.g., spreadsheet software such as Microsoft Excel™, Corel Quattro Pro™, or database programs such as Microsoft Access™ or Paradox™) can be adapted to the present invention by inputting data which is loaded into the memory of a digital system, and performing an operation as noted herein on the data. For example, systems can include the foregoing software having the appropriate genotypic information, associations between phenotype and genotype, etc., e.g., used in conjunction with a user interface (e.g., a GUI in a standard operating system such as a Windows, Macintosh or LINUX system) to perform any analysis noted herein, or simply to acquire data (e.g., in a spreadsheet) to be used in the methods herein.

Systems typically include, e.g., a digital computer with software for performing association analysis and/or risk prediction, as well as data sets entered into the software system comprising genotypes for a set of genetic markers, phenotypic values, and/or the like. The computer can be, e.g., a PC (Intel x86 or Pentium chip- compatible DOS,™ OS2,™ WINDOWS,™ WINDOWS NT,™ WINDOWS95,™ WINDOWS98,™ LINUX, Apple-compatible, MACINTOSH™ compatible, Power PC compatible, or a UNIX compatible (e.g., SUN™ work station) machine) or other commercially common computer which is known to one of skill. Software for performing association analysis and/or risk prediction can be constructed by one of skill using a standard programming language such as Visualbasic, Fortran, Basic, Java, or the like, according to the methods herein.

Any system controller or computer optionally includes a monitor which can include, e.g., a cathode ray tube ("CRT") display, a flat panel display (e.g., active matrix liquid crystal display, liquid crystal display), or others. Computer circuitry is often placed in a box which includes numerous integrated circuit chips, such as a microprocessor, memory, interface circuits, and others. The box also optionally includes a hard disk drive, a floppy disk drive, a high capacity removable drive such as a writeable CD-ROM, and other common peripheral elements. Inputting devices such as a keyboard or mouse optionally provide for input from a user and for user selection of genetic marker genotype, phenotypic value, or the like in the relevant computer system.

- 42 -

The computer typically includes appropriate software for receiving user instructions, either in the form of user input into a set of parameter fields, e.g., in a GUI, or in the form of preprogrammed instructions, e.g., preprogrammed for a variety of different specific operations. The software then converts these instructions to appropriate language for instructing the system to carry out any desired operation. For example, in addition to performing risk prediction, a digital system can control equipment for detecting polymorphisms according to the relevant method herein.

The invention can also be embodied within the circuitry of an application specific integrated circuit (ASIC) or programmable logic device (PLD). In such a case, the invention is embodied in a computer readable descriptor language that can be used to create an ASIC or PLD. The invention can also be embodied within the circuitry or logic processors of a variety of other digital apparatus, such as PDAs, laptop computer systems, displays, image editing equipment, etc.

MOLECULAR BIOLOGICAL TECHNIQUES

Conventional techniques of molecular biology and nucleic acid chemistry, which are within the skill of the art, are fully explained in the literature. See, for example, Sambrook *et al.*, 1989, *Molecular Cloning - A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York; Oligonucleotide Synthesis (M.J. Gait, ed., 1984); *Nucleic Acid Hybridization* (B.D. Hames and S.J. Higgins. eds., 1984); the series, *Methods in Enzymology* (Academic Press, Inc.); the series *Current Protocols in Human Genetics* (Dracopoli *et al.*, eds., 1984 with quarterly updates, John Wiley & Sons, Inc.); Berger and Kimmel, *Guide to Molecular Cloning Techniques*, *Methods in Enzymology* volume 152 Academic Press, Inc., San Diego, CA; Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual* (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 2000; *Current Protocols in Molecular Biology*, F.M. Ausubel *et al.*, eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2004); Freshney (1994) *Culture of Animal Cells*, a *Manual of Basic Technique*, third edition, Wiley- Liss, New York and the references cited therein; Payne *et al.* (1992) *Plant Cell and Tissue Culture in Liquid Systems* John Wiley & Sons, Inc. New York, NY; Gamborg and Phillips (Eds.) (1995) *Plant Cell, Tissue and Organ Culture*; *Fundamental Methods Springer Lab Manual*, Springer-Verlag (Berlin Heidelberg New York); and Atlas and Parks (Eds.) *The Handbook of Microbiological Media* (1993) CRC Press, Boca Raton, FL, all of which are

incorporated herein by reference. All patents, patent applications, and publications mentioned herein, both supra and infra, are incorporated herein by reference.

EXAMPLES

- 5 The following examples are offered to illustrate, but not to limit, the claimed invention. It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and
10 purview of this application and scope of the appended claims.

Example 1: Genotyping Protocol: Allele-specific PCR of FRZB Alleles

- This example describes a method of genotyping SNPs (and/or determining the frequency of alleles in pooled DNA samples) in the FRZB locus, e.g., SNPs that are associated with obesity and/or osteoporosis. The method used was basically that of
5 Germer, et al., *Genome Research*, 10:258-266 (2000). The method and primers were initially tested on cell lines listed in Table 2. The genotypes of these cell lines in FRZB had been previously determined by DNA sequencing for certain of the lines; other lines were genotyped by sequencing later, and some have not been sequenced to date.

- Allele-specific primers and common primers were designed and tested. The allele
10 specific PCR was performed using pairs of the primers listed in Table 2 (SEQ ID NOS:3-29). All primers are shown in the 5' to 3' orientation. In addition to the primers, the control genotype of each cell line control for particular SNPs used in the test assays are also shown in Table 2.

Table 2: Allele-specific Assay Information

SNP	Cell Line Control	Control Genotype	Primer	Allele	Primer Sequences (SEQ ID NO:)
FRZB_C18679T	GM15890	A1/A1	AS1	C	ATAGTAGGAGGAGTACTGTGTCG (SEQ ID NO:3)
FRZB_C18679T	GM15892	A1/A2	AS2	T	AATAGTAGGAGGAGTACTGTGTCA (SEQ ID NO:4)
FRZB_C18679T	GM15891	A2/A2	Common		CCCTGTGGACTATCACCTAATGTT (SEQ ID NO:5)

SNP	Cell Line Control	Control Genotype	Primer	Allele	Primer Sequences (SEQ ID NO:)
FRZB_G19524A	GM15890	A1/A1	AS1	G	ATAGGCCATCAGTTGTGC (SEQ ID NO:6)
FRZB_G19524A	GM15208	A1/A2	AS2	A	CATAGGCCATCAGTTGTGT (SEQ ID NO:7)
FRZB_G19524A	GM15206	A2/A2	Common		TCAAAGTGCTCTGCTGTTTAC (SEQ ID NO:8)
FRZB_T22242A	GM15207	A1/A1	AS1	T	GATCTATTTCTCCATCTGCT (SEQ ID NO:9)
FRZB_T22242A	GM15889	A1/A2	AS2	A	ATCTTATTTCTCCATCTGCA (SEQ ID NO:10)
FRZB_T22242A	GM15206	A2/A2	Common		CGTGAGGGAAAGGAATGTTG (SEQ ID NO:11)
FRZB_G23043A	GM15888	A1/A1	AS1	G	TTGTCTTTTATCCCAGTCATTC (SEQ ID NO:12)
FRZB_G23043A	GM15890	A1/A2	AS2	A	TTGTCTTTTATCCCAGTCATTT (SEQ ID NO:13)
FRZB_G23043A	GM13626	A1/A2	Common		CATCATGGCACTTAGTCTTTATCTC (SEQ ID NO:14)
FRZB_G23415A	GM15888	A1/A1	AS1	G	AAAAATGTAAACCTATAAACTACACG (SEQ ID NO:15)
FRZB_G23415A	GM15890	A1/A2	AS2	A	GAAAAATGTAAACCTATAAACTACACA (SEQ ID NO:16)
FRZB_G23415A	GM13626	A1/A2	Common		TCTTGATTTTATATATGGAATGGGT (SEQ ID NO:17)
FRZB_T23549C	GM15204	A1/A1	AS1	T	ACAGTACTTGAACAAGAAAGACTTAT (SEQ ID NO:18)
FRZB_T23549C	GM15889	A1/A2	AS2	C	ACAGTACTTGAACAAGAAAGACTTAC (SEQ ID NO:19)
FRZB_T23549C	NA14700	A1/A2	Common		ACTGTTCTAAATCTTAGCTGTCCTATTC (SEQ ID NO:20)
FRZB_G24791A	GM10347	A1/A1	AS1	G	CTCCCTTTTGACAAATCTACTG (SEQ ID NO:21)
FRZB_G24791A	GM11235	A1/A2	AS2	A	TCTCCCTTTTGACAAATCTACTA (SEQ ID NO:22)
FRZB_G24791A	GM13625	A2/A2	Common		GAAACTACCCTCCAGTAAGTTCTTC (SEQ ID NO:23)

SNP	Cell Line Control	Control Genotype	Primer	Allele	Primer Sequences (SEQ ID NO:)
					NO:23)
FRZB_C26794G	GM11318	A1/A1	AS1	C	TTCGGGATTAGTTGCG (SEQ ID NO:24)
FRZB_C26794G	GM10873	A1/A2	AS2	G	TTCGGGATTAGTTGCC (SEQ ID NO:25)
FRZB_C26794G	NA14683	A2/A2	Common		GTCTGGCAGGAAC TCGAACC (SEQ ID NO:26)
FRZB_G27014A	GM05045	A1/A1	AS1	G	TGGGGGCAGACTCTTAAG (SEQ ID NO:27)
FRZB_G27014A	GM10873	A1/A2	AS2	A	TGGGGGCAGACTCTTAAA (SEQ ID NO:28)
FRZB_G27014A	NA14683	A1/A2	Common		CATGATTAGTGAAATAGAAAATCACA (SEQ ID NO:29)
FRZB_T19575G	GM10347	A1/A1	AS1	T	GACTGAAGAAGTCAAGTTTGAGT (SEQ ID NO:30)
FRZB_T19575G	GM04340	A1/A2	AS2	G	ACTGAAGAAGTCAAGTTTGAGG (SEQ ID NO:31)
FRZB_T19575G	GM13626	A1/A2	Common		TGAACAGCAGAGCACITTGAT (SEQ ID NO:32)
FRZB_T2303723	GM12548	A1/A1	AS1	T	GTCGGCATTCTTATCATTCA (SEQ ID NO:33)
FRZB_T2303723 C	GM14663	A1/A2	AS2	C	CGGCATTCTTATCATTCG (SEQ ID NO:34)
FRZB_T2303723 C	GM14667	A2/A2	Common		AATAAGTCTCATCCATACTCAACCC (SEQ ID NO:35)

The PCR amplification was carried out in a total reaction volume of 50 µl containing the following reagents:

3.5 ng purified human genomic DNA

5 0.2 µM each primer (one common primer and one allele-specific primer)

50 µM each dATP, dCTP, dGTP

25 µM each dTTP

75 µM each dUTP

10 mM Tris-HCl, pH 8.3

- 46 -

3 mM MgCl₂

0.02U UNG (Uracil-n-glycosylase)

4% DMSO

2% glycerol

5 0.2X SYBR™ Green

12 units CEA2 Gold™ DNA polymerase*

* developed and manufactured by Hoffmann-La Roche. 0.5% glycerol added with CEA2 Gold. 1% DMSO added with SYBR Green.

10 The PCR was run on the GeneAmp 5700 Sequence Detection System (ABI) measuring SYBR™ Green I (Molecular Probes, Eugene, OR, USA) fluorescence in realtime (Higuchi, R. et al., 1993, *Biotechnology* 11:1026-30), as follows:

50°C for 2min

95°C for 12min

95°C for 20 sec

15 58°C for 20 sec

repeated for 45 cycles.

20 All of the primers shown in Table 2 used under these conditions resulted in the correct, known genotype for each SNP for the designated cell line. Thus, the genotyping assay is usable for further analysis of the SNPs. The cell lines were used as positive controls in further analyses.

Example 2: Pooling Analysis for Identifying Disease Association with Known FRZB SNPs

25 Pooling analysis was used to facilitate the screening of candidate genes (e.g., the FRZB gene) thought to be associated with osteoporosis and/or obesity. A large allele frequency difference between pools of DNA from patients and from controls is indicative of a possible involvement of a gene in these conditions. Instead of genotyping a very large number of samples, measuring the allele frequency of a single pool composed of equal amounts of those samples allows the rapid survey of a large number of candidate genes.

Eleven exemplary SNPs in the human FRZB gene are listed in Table 1. The allele frequencies of these SNPs were measured using allele-specific, quantitative PCR on pooled samples from the Study of Osteoporotic Fractures (SOF), using the method of Germer, et al., *Genome Research*, 10:258-266 (2000) as described in Example 1. The
5 Study of Osteoporotic Fractures (Kado, et al., *Arch. Intern. Med.* 159:1215-1220 (1999)) includes DNAs obtained from women age 65 or older exhibiting hip fractures, vertebral fractures, low bone mineral density (BMD) or high body mass index (BMI), as well as control samples. The study group comprises 1042 DNA samples altogether.

Allele Frequency Determination:

10 To measure a SNP allele frequency in a mixture of DNAs pooled from individual samples, equal aliquots of the pool are divided between two PCR reactions, each of which contains a primer pair specific to one or the other SNP allelic variant (e.g., one allele-specific primer and one common primer for each SNP, as described in Example
15 1). The specificity of the PCR amplification is conferred by placing the 3' end of one of the primers (the allele-specific primer) directly over and matching one or the other of the variant nucleotides. This specificity can be enhanced particularly by using the Stoffel fragment of Taq DNA polymerase or variants thereof. Ideally, only completely matched primers are extended, and only the matching allele is amplified. In practice, however,
20 there will typically be amplification of the mismatched allele, but this will occur much less efficiently such that many more amplification cycles are needed to generate detectable levels of product. Mismatch amplification is frequently delayed by >10 cycles when amplification is monitored on a cycle-by-cycle basis using fluorescent dsDNA binding dyes such as SYBR Green I. A delay of around six cycles is adequate for the
25 determination of allele frequencies of SNPs for which the frequency of the minor allele is greater than a few percent.

When the allele frequency is 50%, one expects that each of the two PCR amplifications will require the same number of cycles to produce the same fluorescent signal, assuming that both allele-specific primers amplify with equal efficiency. The number of cycles
30 before a reaction crosses a predetermined threshold, the C_t , can be fractional. When one allele is more frequent, the amplification of that allele will reach the threshold at an earlier cycle, that is, have a smaller C_t . The difference in C_t 's between the two PCR reactions, the ΔC_t , is a measure of the bias and thus of the allele frequency. A one-cycle delay means that the ratio of the amount of one allele to the other is 1:2, a two-cycle

delay, 1:4, or in general, $1:2^{\Delta C_t}$. Converting a ratio to a frequency by adding the numerator to the denominator results in the equation

$$\text{Frequency of allele}_1 = 1/(2^{\Delta C_t} + 1),$$

Where $\Delta C_t = (C_t \text{ of allele}_1\text{-specific PCR}) - (C_t \text{ of allele}_2\text{-specific PCR})$.

- 5 Note that ΔC_t can be either positive or negative, depending on which specific PCR exhibits the lowest C_t . The “2” in the denominator is properly “1 + the initial replication efficiency”. However, the initial replication efficiency is usually close to 100% so that “2” is an adequate approximation. The amplification efficiencies for the two allele-specific PCRs may differ slightly. As discussed in Germer *et al.* (*supra*), this
- 10 can be measured and compensated for by performing the assay on a DNA known to be heterozygous for the SNP of interest. The ΔC_t for this DNA should equal zero if the PCRs are equally efficient. Any deviation from zero indicates that they are not. This deviation can then be subtracted from all ΔC_t measurements to compensate for differential amplification efficiencies.
- 15 Each SNP except T2303723C was described by its position in the reference GenBank accession sequence NT_005100.3. This sequence (containing only 4 exons) was archived and replaced in GenBank with the more complete sequence containing 6 exons (NT_005265), which was in turn replaced by NT_005403. However, the original sequence for which the numbering was based (NT_005100.3) can be obtained from
- 20 GenBank, and the first 30,000 nucleotides of this sequence are presented as SEQ ID NO:1. Thus, for example, the second SNP listed in Table 1 is found at position 18679 of NT_005100.3 (and of SEQ ID NO:1), where a “T” nucleotide is present as the complement of nucleotide 18679 of SEQ ID NO:1. The common allele has a “C” nucleotide at this position. SNP T2303723C was described by its position in the
- 25 GenBank accession sequence NT_005265. The SNPs will be referred to by the SNP # (i.e., by nucleotide position in either SEQ ID NO:1 or NT_005265, as indicated above, although the nucleotide(s) indicated occupy either the indicated position or its complement, depending on the particular SNP; see Figure 1) in the subsequent text. Similarly, the SNPs can be located in any FRZB sequence by performing a sequence
- 30 alignment with the allele-specific primer sequences listed in Table 2. SNPs can also be unambiguously located in the NCBI dbSNP database (<http://www.ncbi.nlm.nih.gov/SNP/index.html>) through the SNP source number listed in Table 1.

- The pooled samples were designated A through D (see, e.g., Table 3). The criterion for pool A was hip fracture (any incident hip fracture since baseline, excluding prior hip fracture since age 50). The criterion for pool B was vertebral fracture (incident vertebral fracture by morphometry between baseline and visit 3). The criterion for pool C was low BMD (bone mineral density, defined as having hip BMD T-score <-2.5). The criteria for pool D (the control) were no fracture since age 50 and hip BMD Z-score >1.285 . The criterion for the High BMI pool was a body mass index in the highest 5% of SOF participants and the criterion for the Low BMI pool was a body mass index in the lowest 5% of SOF participants.
- The pools involved combining equal amounts of DNAs from patients with each specific criterion. Pool A included 275 patient samples; pool B contained 262 patient samples; pool C included 276 patient samples; and pool D contained 278 patient samples. The High BMI pool contained 141 samples and the Low BMI pool contained 82 samples. The PCRs were performed in 4 replicates using the primers and protocol outlined in Example 1 except that the PCRs contained 10 ng purified human genomic DNA and 2 μ M Rox. The averages of the four replicates were calculated, and the allele frequencies were then calculated following the method above and compared to the controls. A change in allele frequency of greater than approximately 4-5% was considered significant. The pooling results for each FRZB SNP are shown in Table 3 below.

Table 3: FRZB Pooling results

Pool	Average Allele 1 Frequency (%)						Change in Allele 1 Frequency (%)			
	A	B	C	D	Low BMI	High BMI	A-D	B-D	C-D	Low BMI-High BMI
FRZB_C18679T	66.38	61.81	69.06	63.79	71.02	65.75	2.59	-1.98	5.27	5.27
FRZB_G19524A	74.12	68.57	71.49	68.6	81.69	67.6	5.52	-0.03	2.89	14.09
FRZB_T22242A	74.66	69.91	72.74	70.14	82.27	72.18	4.52	-0.23	2.60	10.09
FRZB_G23043A	88.01	91.66	91.15	88.65	88.79	88.89	-0.63	3.01	2.50	-0.10
FRZB_G23415A	83.69	89.48	88.86	84.90	85.30	86.26	-1.21	4.58	3.96	-0.96
FRZB_T23549C	98.30	97.89	97.33	98.56	97.77	97.02	-0.26	-0.67	-1.22	0.75
FRZB_A24791G	50.53	56.92	55.67	56.26	46.12	61.57	-5.73	0.66	-0.59	-15.45
FRZB_C26794G	82.22	81.82	78.24	82.23	80.5	76.66	0.0	-0.41	-3.99	3.84
FRZB_G27014A	96.79	98.69	96.54	95.5	98.62	96.1	1.29	3.2	1.04	2.52

- A Hip Fracture
 B Vertebral Fracture
 C Low BMD
 D Control

Example 3: Individual Genotyping

Because of the significant differences in allele frequency between some osteoporotic case and control groups and between the high and low BMI groups, the individual samples were genotyped to both verify the allele frequency differences and to determine genotype frequencies in these groups. The same allele-specific protocol was used on individual DNA samples to determine their genotypes. The PCR amplifications were performed as described in Example 1. The primers used to genotype the FRZB SNPs were those listed in Table 2. These results were analyzed using Pearson's Chi-Square (or an Exact Test if the number of subjects with one of the genotypes was 10 or less) to determine whether the distribution of genotype frequencies were significantly different between the groups of individuals who had any of the osteoporotic phenotypes and controls and between the groups of individuals with high BMI (a.k.a., obesity) and low BMI.

A significant association between an increased BMI and the FRZB_T2303723C, FRZB_C18679T, FRZB_G19524A and FRZB_T22242A SNPs was demonstrated ($p < 0.05$). A lesser association between an increased BMI and the FRZB_A24791G SNP was also demonstrated ($p < 0.1$). The T allele of T2303723C, the T allele of C18679T, the A allele of G19524A, the A allele of T22242A, and the G allele of A24791G were each demonstrated to be associated with increased BMI (and thus, with an increased risk of obesity). The association is most statistically significant if the effect of each of these alleles is assumed to be recessive.

With respect to osteoporosis, significant associations were demonstrated between increased incidence of vertebral fracture and the FRZB_C26794G and FRZB_G27014A SNPs ($p < 0.05$). A lesser association between increased incidence of hip fracture and the FRZB_C18679T SNP was also demonstrated ($p < 0.1$), as well as between increased incidence of hip fracture and the FRZB_G19524A SNP ($p < 0.1$). The C allele of C18679T, the G allele of G19524A, the C allele of C26794G, and the G allele of G27014A were each demonstrated to be associated with increased incidence of hip or vertebral fracture (and thus, with an increased risk of osteoporosis). The association is most statistically significant if the effect of each of these alleles is assumed to be recessive.

In addition, it is worth noting that after adjustment for weight and age of the patients, the A allele of A24791G is associated with an increased incidence of hip fracture ($p < 0.1$; with the A allele assumed to be recessive). Similarly, using an alternative control group (defined as no fracture since age 50 and highest 5% hip BMD by individual 5-year age

group), the T allele of T22242A is associated with an increased incidence of hip fracture. Again, the association is most statistically significant if the effect of this allele is assumed to be recessive.

Association of FRZB SNPs with Obesity in Women

- 5 This example demonstrates the association of FRZB SNPs with obesity in women.

As noted, FRZB genotyping was carried out on women from the Study of Osteoporotic Fractures (SOF), using a genotyping method essentially as described in Example 1. Table 4 lists the genotyping results for the four FRZB SNPs indicated and the numbers and percents of individuals in each BMI category for each genotype.

10

Table 4: FRZB genotype association with BMI

	C18679T					G19524A			Overall Distribution of G19524A P=.0051
	BMI	No.	%			BMI	No.	%	
T/T	High	18	12.8		A/A	High	14	10.0	
	Low	3	3.7			Low	0	0.0	
T/C	High	55	39.0		A/G	High	50	35.5	Model where 19524A is recessive allele P=.0032
	Low	38	46.3			Low	30	36.6	
C/C	High	68	48.2		G/G	High	77	54.6	
	Low	41	50.0			Low	52	63.4	
	Total No. high BMI=141; Total No. low BMI=82								
	T22242A					A24791G			
	BMI	No.	%.			BMI	No.	%	
A/A	High	15	10.7		G/G	High	36	25.5	
	Low	2	2.4			Low	13	15.9	
A/T	High	57	40.4		G/A	High	69	48.9	
	Low	32	39.0			Low	43	52.4	
T/T	High	69	48.9		A/A	High	36	25.5	
	Low	48	58.5			Low	26	31.7	

Statistical analysis, methods and algorithms: Association of FRZB genotypes with obesity was assessed using Pearson's Chi-Square test or, if the number of subjects with one of the genotypes was 10 or less, Fisher's Exact Test. Table 4 shows the p-values

(probabilities) that the distribution of genotypes at the G19524A SNP between the high and low BMI groups could have been obtained purely by chance. This is seen to be highly unlikely. Thus there is a statistical association between G19524A and BMI. The p-value is even lower if it is assumed that the minor allele, 19524A, exerts its genetic effect in a recessive (two 19524A alleles required) mode.

Association of FRZB With Osteoporosis in SOF Samples

This example demonstrates the association of FRZB SNPs with osteoporosis in SOF samples.

Table 5 lists the genotyping results for FRZB SNP C26794G and the numbers and percents of individuals in the vertebral fracture and control category for each genotype.

Table 5: FRZB genotype association with vertebral fracture

	C26794G (Arg -> Gly)			Overall Distribution
	fracture	No.	%	
G/G	No	1	0.4	P=.025
	Yes	2	0.8	
G/C	No	52	18.7	
	Yes	29	11.1	
C/C	No	225	80.9	
	Yes	231	88.1	

Total No. vertebral fracture =262; Total No. control =278

Statistical analysis, methods and algorithms: Association of FRZB genotypes with osteoporosis was assessed using Pearson's Chi-Square test or, if the number of subjects with one of the genotypes was 10 or less, Fisher's Exact Test. Table 5 shows the p-value (probability) that the distribution of genotypes at C26794G between the vertebral fracture and control groups could have been obtained purely by chance. This is seen to be highly unlikely. Thus there is a statistical association between C26794G and vertebral fracture. C26794G imparts a coding (amino acid) substitution (arginine to glycine) upon the protein sequence of FRZB.

As FRZB is a small component of the complex system of genes associated with obesity and/or osteoporosis (which are thought to be related since bone and fat cells originate from branches along the same developmental pathway), the effect of the FRZB locus is expected to be variable. Other factors, such as eating habits, exercise, diet, general health and the presence of associated diseases, may exert dominating effects which, in some cases, may mask the effect of the FRZB genotypes. Furthermore, because allele frequencies at other loci relevant to weight and bone-related diseases differ between populations and, thus, populations exhibit different risks for such diseases, it is expected that the effect of the FRZB genotype may be of different magnitude in some populations. Although the contribution of the FRZB genotype may, in certain populations, be relatively minor by itself, genotyping at the FRZB locus will contribute information that is, nevertheless, useful for a characterization of an individual's predisposition towards obesity and/or osteoporosis. The FRZB genotype information may be particularly useful when combined with genotype information from other loci and/or clinical tests for obesity and/or osteoporosis, for example.

Although the invention has been disclosed in the context of certain embodiments and examples, it will be understood by those skilled in the art that the invention extends beyond the specifically disclosed embodiments to other alternative embodiments and/or uses and obvious modifications and equivalents thereof. Accordingly, the invention is not intended to be limited by the specific disclosures of preferred embodiments herein, but instead by reference to claims attached hereto. All references cited herein are hereby incorporated by reference in their entireties.

While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. For example, all the techniques, apparatus and/or compositions described above can be used in various combinations. All publications, patents, patent applications, and/or other documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, and/or other document were individually indicated to be incorporated by reference for all purposes.

CLAIMS

WHAT IS CLAIMED IS:

1. A method for determining an individual's risk for obesity, the method
5 comprising: detecting presence of at least one obesity-related polymorphism in a frizzled-related protein (FRZB) gene in a nucleic acid sample of the individual, wherein the presence of said at least one polymorphism provides an indication of the individual's risk for obesity.
2. The method of claim 1, wherein the individual's risk for obesity is an increased
10 risk as compared to an individual without the at least one polymorphism.
3. The method of claim 1, wherein the individual's risk for obesity is a decreased risk as compared to an individual without the at least one polymorphism.
4. The method of claim 1, wherein the at least one polymorphism comprises a predisposing or protective polymorphism in the FRZB gene.
- 15 5. The method of claim 1, wherein the nucleic acid sample comprises DNA or RNA.
6. The method of claim 1, wherein the at least one polymorphism comprises at least one single nucleotide polymorphism (SNP).
7. The method of claim 6, wherein the at least one polymorphism is selected from
20 the group consisting of: T allele of T2303723C, C allele of T2303723C, C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, T allele of T22242A, A allele of T22242A, A allele of A24791G, and G allele of A24791G.
8. The method of claim 1, wherein the at least one polymorphism comprises two or more polymorphisms.
- 25 9. The method of claim 8, wherein at least one of the two or more polymorphisms is selected from the group consisting of: T allele of T2303723C, C allele of T2303723C, C allele of C18679T, T allele of C18679T, G allele of G19524A, A

- 55 -

allele of G19524A, T allele of T22242A, A allele of T22242A, A allele of A24791G, and G allele of A24791G.

10. The method of claim 8, wherein the two or more polymorphisms are selected from the group consisting of: T allele of T2303723C, C allele of T2303723C, C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, T allele of T22242A, A allele of T22242A, A allele of A24791G, and G allele of A24791G.
11. The method of claim 1, wherein the at least one polymorphism is detected by sequencing.
- 10 12. The method of claim 1, wherein the at least one polymorphism is detected by amplification.
13. The method of claim 12, wherein the amplification comprises a polymerase chain reaction or a ligase chain reaction.
14. The method of claim 1, wherein the detecting comprises:
 - 15 - contacting the nucleic acid sample with at least one sequence-specific oligonucleotide under conditions that allow binding of said at least one oligonucleotide to the nucleic acid sample, wherein the at least one sequence-specific oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the at least one obesity-related polymorphism; and,
 - 20 - detecting the hybridization of the at least one oligonucleotide to the nucleic acid sample.
15. The method of claim 1, wherein the detecting comprises:
 - 25 - amplifying the nucleic acid sample, thereby providing an amplified nucleic acid sample;
 - contacting the amplified nucleic acid sample with at least one sequence-specific oligonucleotide under conditions that allow binding of the oligonucleotide to the amplified nucleic acid sample,

wherein the at least one sequence-specific oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the at least one obesity-related polymorphism; and,

5 - detecting the hybridization of the at least one sequence-specific oligonucleotide to the amplified nucleic acid sample.

16. The method of claim 1, wherein detecting the presence of the at least one obesity-related polymorphism comprises qualitatively detecting the presence of the at least one obesity-related polymorphism.
17. The method of claim 1, wherein detecting the presence of the at least one
10 obesity-related polymorphism comprises quantitatively detecting the presence of the at least one obesity-related polymorphism.
18. The method of claim 1, wherein the presence of the polymorphism inherited from one of the individual's parents provides an indication of the individual's risk for obesity, or wherein the presence of the polymorphism inherited from
15 both of the individual's parents provides an indication of the individual's risk for obesity.
19. The method of claim 1, comprising performing at least one clinical test for obesity.
20. The method of claim 19, wherein performing the at least one clinical test for
20 obesity comprises determining a body mass index (BMI) of the individual.
21. A method for determining an individual's risk for osteoporosis, the method comprising: detecting presence of at least one osteoporosis-related polymorphism in an FRZB gene in a nucleic acid sample of the individual, wherein the presence of said at least one polymorphism provides an indication of
25 the individual's risk for osteoporosis.
22. The method of claim 21, wherein the individual's risk for osteoporosis is an increased risk as compared to an individual without the at least one polymorphism.

23. The method of claim 21, wherein the individual's risk for osteoporosis is a decreased risk as compared to an individual without the at least one polymorphism.
24. The method of claim 21, wherein the at least one polymorphism comprises a predisposing or protective polymorphism in the FRZB gene.
25. The method of claim 21, wherein the nucleic acid sample comprises DNA or RNA.
26. The method of claim 21, wherein the at least one polymorphism comprises at least one single nucleotide polymorphism (SNP).
27. The method of claim 26, wherein the at least one polymorphism is selected from the group consisting of: C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, A allele of A24791G, G allele of A24791G, C allele of C26794G, G allele of C26794G, G allele of G27014A, and A allele of G27014A.
28. The method of claim 21, wherein the at least one polymorphism comprises two or more polymorphisms.
29. The method of claim 28, wherein at least one of the two or more polymorphisms is selected from the group consisting of: C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, A allele of A24791G, G allele of A24791G, C allele of C26794G, G allele of C26794G, G allele of G27014A, and A allele of G27014A.
30. The method of claim 28, wherein the two or more polymorphisms are selected from the group consisting of: C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, A allele of A24791G, G allele of A24791G, C allele of C26794G, G allele of C26794G, G allele of G27014A, and A allele of G27014A.
31. The method of claim 21, wherein the at least one polymorphism is detected by sequencing.

32. The method of claim 21, wherein the at least one polymorphism is detected by amplification.
33. The method of claim 32, wherein the amplification comprises a polymerase chain reaction or a ligase chain reaction.
- 5 34. The method of claim 21, wherein the detecting comprises:
- contacting the nucleic acid sample with at least one sequence-specific oligonucleotide under conditions that allow binding of the oligonucleotide to the nucleic acid sample, wherein the at least one sequence-specific oligonucleotide hybridizes under stringent
10 conditions to a region of the FRZB gene comprising the at least one osteoporosis-related polymorphism; and,
 - detecting the hybridization of the at least one sequence-specific oligonucleotide to the nucleic acid sample.
35. The method of claim 21, wherein the detecting comprises:
- 15 - amplifying the nucleic acid sample, thereby providing an amplified nucleic acid sample;
 - contacting the amplified nucleic acid sample with at least one sequence-specific oligonucleotide under conditions that allow
20 binding of the oligonucleotide to the amplified nucleic acid sample, wherein the at least one sequence-specific oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the at least one osteoporosis-related polymorphism; and,
 - detecting the hybridization of the at least one sequence-specific oligonucleotide to the amplified nucleic acid sample.
- 25 36. The method of claim 21, wherein detecting the presence of the at least one osteoporosis-related polymorphism comprises qualitatively detecting the presence of the at least one osteoporosis-related polymorphism.

37. The method of claim 21, wherein detecting the presence of the at least one osteoporosis-related polymorphism comprises quantitatively detecting the presence of the at least one osteoporosis-related polymorphism.
- 5 38. The method of claim 21, wherein the presence of the polymorphism inherited from one of the individual's parents provides an indication of the individual's risk for osteoporosis, or wherein the presence of the polymorphism inherited from both of the individual's parents provides an indication of the individual's risk for osteoporosis.
- 10 39. The method of claim 21, comprising performing at least one clinical test for osteoporosis.
40. The method of claim 39, wherein the at least one clinical test for osteoporosis comprises a bone-turnover assay or a bone scan.
- 15 41. A method for determining an individual's risk for obesity and/or osteoporosis, the method comprising: determining the individual's genotype at one or more polymorphic sites in an FRZB gene, wherein a first genotype at the one or more polymorphic sites is statistically associated with an increased risk for obesity and/or osteoporosis as compared to a second genotype at the one or more polymorphic sites.
- 20 42. The method of claim 41, wherein at least one of the one or more polymorphic sites consists of a single nucleotide position.
43. The method of claim 42, wherein the individual's genotype is determined at a plurality of polymorphic sites in the FRZB gene, and wherein each of the polymorphic sites consists of a single nucleotide position.
- 25 44. The method of claim 41, wherein at least one of the one or more polymorphic sites is selected from the group consisting of: nucleotide position 2628, nucleotide position 18679, nucleotide position 19524, nucleotide position 22242, nucleotide position 24791, nucleotide position 26794, and nucleotide position 27014 of SEQ ID NO:1.

45. The method of claim 41, wherein the first genotype is statistically associated with an increased risk for obesity as compared to the second genotype; and wherein the first genotype comprises two T alleles and the second genotype two C alleles or one T allele and one C allele of SNP T2303723C, the first genotype comprises
5 two T alleles and the second genotype two C alleles or one T allele and one C allele of SNP C18679T, the first genotype comprises two A alleles and the second genotype two G alleles or one A allele and one G allele of SNP G19524A, the first genotype comprises two A alleles and the second genotype two T alleles or one A allele and one T allele of SNP T22242A, and/or the first genotype comprises two
10 G alleles and the second genotype two A alleles or one G allele and one A allele of SNP A24791G.
46. The method of claim 41, wherein the first genotype is statistically associated with an increased risk for osteoporosis as compared to the second genotype; and wherein the first genotype comprises two C alleles and the second genotype two
15 T alleles or one T allele and one C allele of SNP C18679T, the first genotype comprises two G alleles and the second genotype two A alleles or one A allele and one G allele of SNP G19524A, the first genotype comprises two A alleles and the second genotype two G alleles or one G allele and one A allele of SNP A24791G, the first genotype comprises two C alleles and the second genotype two G alleles
20 or one C allele and one G allele of SNP C26794G, and/or the first genotype comprises two G alleles and the second genotype two A alleles or one G allele and one A allele of SNP G27014A.
47. The method of claim 41, wherein determining the individual's genotype comprises obtaining a nucleic acid sample from the individual.
- 25 48. The method of claim 47, wherein determining the individual's genotype comprises amplifying at least a portion of the FRZB gene from the nucleic acid sample, the portion comprising at least one of the one or more polymorphic sites.
- 30 49. The method of claim 47, wherein determining the individual's genotype comprises performing an allele-specific amplification or an allele-specific extension reaction.

50. The method of claim 47, wherein determining the individual's genotype comprises sequencing at least a portion of the FRZB gene from the nucleic acid sample, the portion comprising at least one of the one or more polymorphic sites.
- 5 51. The method of claim 47, wherein at least one of the one or more polymorphic sites consists of a single nucleotide position, the method comprising:
- contacting the nucleic acid sample with at least one sequence-specific oligonucleotide under stringent conditions, wherein the
10 oligonucleotide hybridizes under the stringent conditions to the nucleic acid sample when a first nucleotide occupies the nucleotide position defining the polymorphic site but not when a second nucleotide occupies the nucleotide position, and
 - detecting hybridization of the oligonucleotide to the nucleic acid sample.
- 15 52. A kit for detecting presence of a first predisposing or protective polymorphism in an FRZB gene in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed, the kit comprising: one or more first oligonucleotides capable of detecting the first polymorphism, and
20 instructions for detecting the first polymorphism with the one or more first oligonucleotides and for correlating said detection to the individual's risk for osteoporosis and/or obesity, packaged in one or more containers.
53. The kit of claim 52, wherein the first polymorphism is a single nucleotide polymorphism.
- 25 54. The kit of claim 53, wherein the first polymorphism is selected from the group consisting of: T allele of T2303723C, C allele of T2303723C, C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, T allele of T22242A, A allele of T22242A, A allele of A24791G, G allele of A24791G, C allele of C26794G, G allele of C26794G, G allele of G27014A, and A allele of G27014A.
- 30 55. The kit of claim 52, wherein the one or more first oligonucleotides comprise at least one probe.

56. The kit of claim 52, wherein the first oligonucleotide hybridizes under stringent conditions to a region of the FRZB gene comprising the first polymorphism.
57. The kit of claim 56, wherein the first polymorphism is a first single nucleotide polymorphism comprising a first nucleotide at a first nucleotide position, and
5 wherein, under stringent conditions, the first oligonucleotide hybridizes to a region of the FRZB gene comprising the first single nucleotide polymorphism with a signal to noise ratio that is at least 2x that at which the first oligonucleotide hybridizes to the region of the FRZB gene comprising a second nucleotide at the first nucleotide position.
- 10 58. The kit of claim 56, wherein the first oligonucleotide is fully complementary to the region of the FRZB gene comprising the first polymorphism.
59. The kit of claim 56, wherein the first oligonucleotide comprises at least about 10 contiguous nucleotides complementary to the FRZB gene.
- 15 60. The kit of claim 52, wherein the one or more first oligonucleotides comprise a label.
61. The kit of claim 60, wherein the label comprises an isotopic, fluorescent, fluorogenic or colorimetric label.
62. The kit of claim 60, comprising a reagent, wherein the reagent detects the label.
- 20 63. The kit of claim 52, wherein the one or more first oligonucleotides comprise one or more primers.
64. The kit of claim 63, wherein the first polymorphism is a first single nucleotide polymorphism comprising a first nucleotide at a first nucleotide position, and wherein the 3' nucleotide of one of the one or more first oligonucleotides is complementary to the first nucleotide.
- 25 65. The kit of claim 52, wherein the one or more first oligonucleotides comprise amplification primers, wherein the amplification primers amplify a nucleic acid sequence comprising the first polymorphism.

- 63 -

66. The kit of claim 52, wherein the one or more first oligonucleotides comprise sequencing primers, wherein the sequencing primers flank the first polymorphism.
67. The kit of claim 52, wherein the one or more first oligonucleotides are immobilized on a substrate.
68. The kit of claim 52, comprising one or more second oligonucleotides capable of detecting a second polymorphism.
69. An array for detecting presence of one or more predisposing and/or protective polymorphisms in an FRZB gene in a nucleic acid sample of an individual whose risk for osteoporosis and/or obesity is being assessed, the array comprising:
- a plurality of oligonucleotides, each of which oligonucleotides hybridizes to a region of the FRZB gene comprising at least one of the polymorphisms, whereby said hybridization detects the presence of the polymorphism, whereby said detection provides an indication of the individual's risk for osteoporosis and/or obesity; and,
 - a substrate on which the plurality of oligonucleotides are immobilized.
70. The array of claim 69, wherein the one or more polymorphisms comprise one or more single nucleotide polymorphisms.
71. The array of claim 70, wherein at least one of the one or more polymorphisms is selected from the group consisting of: T allele of T2303723C, C allele of T2303723C, C allele of C18679T, T allele of C18679T, G allele of G19524A, A allele of G19524A, T allele of T22242A, A allele of T22242A, A allele of A24791G, G allele of A24791G, C allele of C26794G, G allele of C26794G, G allele of G27014A, and A allele of G27014A.
72. The array of claim 70, wherein, under stringent conditions, each of the oligonucleotides hybridizes to a region of the FRZB gene comprising one of the single nucleotide polymorphisms with a signal to noise ratio that is at least 2x

that at which the oligonucleotide hybridizes to a region of the FRZB gene comprising any of the remaining single nucleotide polymorphisms.

73. The array of claim 70, wherein each of the oligonucleotides hybridizes to a distinct single nucleotide polymorphism.
- 5 74. The array of claim 69, wherein each of the plurality of oligonucleotides is immobilized at a known, pre-determined position on the substrate.
75. The array of claim 69, wherein each of the plurality of oligonucleotides is fully complementary to a region of the FRZB gene comprising one of the polymorphisms.
- 10 76. The array of claim 69, wherein each of the plurality of oligonucleotides comprises at least about 10 contiguous nucleotides complementary to the FRZB gene.
77. The array of claim 69, wherein each of the plurality of oligonucleotides comprises a label.
- 15 78. A system, comprising: the array of claim 69 and system instructions that correlate said detection of the presence of one or more predisposing or protective polymorphisms to the individual's risk for osteoporosis and/or obesity.

- 1/1 -

5' TGTCGTCTG ACAGCAGAC ₅ '	A	5' TGTCATCTG ACAGTAGAC ₅ '	B
5' TGTGCTTCA ACACGAAGT ₅ '	C	5' TGTGTTTCA ACACAAAGT ₅ '	D
5' AAAGACTCA TTTCTGAGT ₅ '	E	5' AAAGCCTCA TTTCGGAGT ₅ '	F
5' TCAAAGCAG AGTTTCGTC ₅ '	G	5' TCAATGCAG AGTTACGTC ₅ '	H
5' CATTCGGGC GTAAGCCCG ₅ '	I	5' CATTTGGGC GTAAACCCG ₅ '	J
5' TATGCGTGT ATACGCACA ₅ '	K	5' TATGTGTGT ATACACACA ₅ '	L
5' ACATATAAG TGTATATTC ₅ '	M	5' ACATGTAAG TGTACATTC ₅ '	N
5' CCTTTAGTA GGAAATCAT ₅ '	O	5' CCTTCAGTA GGAAGTCAT ₅ '	P
5' AGCAGGCAA TCGTGCGTT ₅ '	Q	5' AGCAGGCAA TCGTCCGTT ₅ '	R
5' TTAAGTATA AATTCATAT ₅ '	S	5' TTAAATATA AATTTATAT ₅ '	T
5' CCCATGAAT GGGTACTTA ₅ '	U	5' CCCACGAAT GGGTGCTTA ₅ '	V

FIG. 1

- 1/24 -

SEQUENCE LISTING

<110> F. Hoffmann-La Roche AG

<120> ASSOCIATIONS OF POLYMORPHISMS IN THE PRZB GENE WITH OBESITY AND
OSTEOPOROSIS

<130> 21634 WO

<150> US 60/464,372

<151> 2003-04-21

<150> US 60/526,689

<151> 2003-12-01

<160> 35

<170> PatentIn version 3.3

<210> 1

<211> 30000

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (7061)..(7160)

<223> n is unidentified (a, c, g or t)

<220>

<221> misc_feature

<222> (7344)..(7344)

<223> n is unidentified (a, c, g or t)

<220>

<221> misc_feature

<222> (7438)..(7438)

<223> n is unidentified (a, c, g or t)

<220>

<221> misc_feature

<222> (7509)..(7608)

<223> n is unidentified (a, c, g or t)

<220>

<221> misc_feature

<222> (28902)..(28902)

<223> n is unidentified (a, c, g or t)

<220>

<221> misc_feature

<222> (28954)..(28954)

<223> n is unidentified (a, c, g or t)

<400> 1

gaattctaga tttgccatta attttgtaca ctgaactagt cgcttgacat tgacattcat 60

tagggattgt gaagaacaaa agaaatagta atatgaagtt actgataaac ggtaaagcac 120

aatataaatg tttaacaaca acgtcattgg ggagaggacg tgtcatcctc aggacaggaa 180

- 2/24 -

ccattagcaa cacaataaaa gcaaccaaca cttatgttct aggtgctgtc ctgagtgc	240
tttgcacatt cattattatt cccatttttg agaaaaggaa gccaaggcat atagagaagc	300
taagttactg tocaagttag cagagctagt aagaaaaaga gaattcgaag cagagcctgt	360
tctacactac actttgtccc ttaagaaatg taattcctaa ctttctagag aatctgttat	420
tctttgccta gcagattgca gttaaaggaa caaaggtttt ctcatgtat tttgaagtac	480
acttatctat ggcttaagag aataaaagggt ggcttctaga caatggattc aaagagataa	540
gagattctga cttcactgca attttgagat agtagagatt atacattcca gattctcatg	600
tccctagtct ctgggtctca gcatgcccc gcgtaactc atcatttccc catagtcatg	660
cacttgtttc tttctagtat tttctatatt gttgaaagggt accaccatac cttagtagtc	720
caggcctggg aggagtcctt tttttttttt tttttttgag acagtctcgc tctgtcactc	780
agactggagt gcagtggcgt gatctctgct cactgcaacc tctgcctcca aggttcaagc	840
aattctcctg tctcagcctc ctgagtagct gggactacag gtgcatgcca ccacgccag	900
ctaatttttg tatttctgtt agagatgggt tttcgccatg ttggccaggc tggctctgaa	960
ctcctgacct cagatgatct gcctgccttg gcctcgcagt gctgagatta cagggtgtgag	1020
tcaccatacc tggctgtggg aagcagtcct gattctacct ttttcctcat tcccctcact	1080
caataggtat cacatgctgt ttggtctact ctgactcaaa ttcccttctc tccctctcta	1140
atgctacttt cttagtttgg accttaccat ttctcaccag gattatggcc ataggatgat	1200
aatggctctc cttatctcca gtatttcccc tccatattcc aatagacatt ttttctgaaa	1260
tgcaaatctt ttgcttaaaa tccttactaa taatatatat tttcatttag ggtaatttgt	1320
tgttagaatt cctaagttta actttctgac aataattttt tgatcaatta ttagaaaatt	1380
tttgttttct gtcttacagt tctaacaatt gacttctaag catttgtaaa agcgtagaaa	1440
cattttcctt tgaataagaa tggaagaagt aaaacactct cttggctta gctattttga	1500
agtgattccc tttaccaag caaagtttta tttttcactc ttatttctac ctttgaacaa	1560
taaacatttc aaatatgggc caatataatt tcttccttta tgaagacatt atttttgtat	1620
gtagtttatc acatgattat tcagaaatag aaaaaccact ggcacgttac tgtaatactt	1680
gggtacattt atcatgaata aatgtttgaa tataatgaaa gtattagtaa ttaatagttt	1740
atgcccaagt tgagccaaat gttttagaat tagtgtttgt tacttgctta tctgtttgtt	1800
tgctgactgc tgtcttttct tggatattgc tttttaaata cttgctgatg attactaaac	1860
ctgggaaagt aattcattaa ttgtaattgt ggggtgtata gtcctaaaat agtattaacc	1920

- 3/24 -

ccttggttaa	tttattcctg	ataggaccta	taaacaaaaa	ggacaaagga	agaatactat	1980
tttctcctag	aataaataga	ggcattaaat	tagtttagca	taaagtgtta	ctacaaatgt	2040
aaaactctta	tataaagtac	taaattggag	tgtcaataag	gatatattgc	taggtacagc	2100
atctctcagc	acggtgttcc	gctggtgcac	tgttattcat	tagaggctgc	tttcttaatt	2160
agaacacaga	aaacctttat	cattactatt	ttctgggatt	atggcatcct	aaaaagaatc	2220
aatagcaatt	tctaattatt	gtcacatcta	ttaattgcac	tggttgaat	agtccttgaa	2280
tagtagggtt	cttttctggt	gaagaattat	ttagcctgca	gttaaaggac	atgagaatga	2340
tttctgagtg	ttcggttagg	aactaaggat	ttaaataat	accagttcag	tgtgatattg	2400
ccatttattg	aagatgcgtt	tggttttttt	tttgtcatca	agtcataagt	gttggtgag	2460
tgcctaatat	ttgcataggc	tttatcatga	taggatgctt	atcatttaca	aattttgtga	2520
aattcaaata	aattattaaa	ttctaaaatt	ttatgcacag	ttgagtattc	agaagagtat	2580
ttccttataa	ccacatgaaa	aaataagtct	catccatact	caacccatga	atgataagaa	2640
tgccgacctc	atgacgttaa	taaaaatcta	cttccttcag	acaattggat	tcttacccaa	2700
atcaaacttg	tgactgagtt	agatttttat	cttttagaat	tcttgaaata	gatattgtat	2760
agttcttttt	tctgtcacct	cctttcagat	tttcctatgg	attctagtaa	cggaaactgt	2820
agaggggcaa	gcagtggtga	gtatatcatt	atcttcctgg	ttttcatttg	tggtcagatt	2880
agtcattgat	gtaatttaga	tacttaagtc	attcatgggg	aagcttgtgc	cctacatttt	2940
acaagtaaac	tttttaaaat	tccctcattt	atattaattt	tatttttaaa	attagcacgt	3000
agtaaaattg	gacttttatt	tggaatata	gtttgtgaca	gatgaatttt	aatattatgt	3060
agagatttat	gtaaccacca	ccacatcagg	atacagaaca	attccatcac	ctcccaaaac	3120
tccttcaagc	tgtatccctt	tataattaca	agttttccct	atcttcatcc	actggcaatc	3180
actcatcttt	tctttatcac	tatagttttc	tctatttgag	gttcgattgt	ccatatagac	3240
agactgacag	taogcctttg	agacttaacc	aattgctgca	tatatcaatg	gtgcattcct	3300
ttttattaca	gaatatttta	ttacttggat	gtatgaccat	ttgttcaccc	attcactcct	3360
tcaaggaagt	ttgggttggt	tccagtttgg	agcaatcgcg	aatagagcta	ctataaacat	3420
tccagtacag	gtttttgtgg	gaatgtaagt	ttttatttct	ctaggctaaa	taagtaggaa	3480
taggatcact	gggttatatg	gtaagtatat	gttttagctt	gtaaaaaagt	gccaaaactgt	3540
ttccaagatg	gctgttcaat	tttgtgctag	taatgtataa	gagttccagt	tgttctatat	3600
ctttgcgagc	ccttggtatt	atcaatatat	tttattgtta	gccattttta	taggtatgtg	3660
ataatagctc	atcatggttt	taaattgcat	ttccctaatt	gctaattgat	ctgaatacat	3720

- 4/24 -

tttcatgtgt ttatttgcta tccacatata tgctttttgg taaaatgttc aataattttg	3780
cccattttta gttgggttgt tttcttaatg agttctgagg gttctttaca tattctggat	3840
acaaaaaaca tcttttggtt ttattcattt tctctattgt ttcactgttt tcaattttat	3900
tgattatttc cttctgcttg cttgcttttag aggtttattt tgctttttct tttgtttctt	3960
agagtagaag cttatattgt gttttgagac tttttttttt tttttgagat ggagtcttgc	4020
tctgttgccc aggtctggagt gccgtgatgc aatctcagct caccacaacc tccacttcct	4080
gggttcaagc aattctgcct cagcctccca actagctagg attacaggtg tgcaaccacca	4140
caccagcta atttttgtat tattagtaga gacaagattt caccatgttg gccagctggg	4200
ctctaact tgatctcagg tgatccaccc accttggcct tacagagtgc tgggattaca	4260
cacgtgagcc accatgtccg gccaaagacct ttcttctaata ataatgtaata gctataaata	4320
cccatctaag aatgacttta gttgcatctc acaaattttg atagtttcat ttttcttcag	4380
ttcaaaatat tttatatttt tcttgatat ttcctttgac ttgtgcatca tttgggatta	4440
tgctgtttta tttgctgctt tttgaggatt ttcttattat ctttgcattt tttcaaattc	4500
tactgttgaa gaatatactt tgtgtgattt cagttctttt agatttatta aggtttattc	4560
tctgatccag gatataatct gtcttggtga atgttccatg tacacttaga aaaaagagta	4620
ttttgctgtt ggggtggaatg tcctttaaat gtcaattcaa tttagttgat taatactgtt	4680
atthagttct tttatattct tgctgatttt ctgtctacta gatctattga ttaccgagag	4740
aagggtgatt aggtctccaa atattgtgga tttttctatt tcttttttca gtctgtcagt	4800
tttgcttcat gtattttgaa gctctgtttg gtacatactt gtttaataatt ttttatgtct	4860
tattggtgaa ttgatgtttt cattatgtag tgtcctttta tccttggtaa ttttctctgc	4920
tctaaattat atttttctg atattgctat aggttaattca gctgtcattt gattagtgtt	4980
tgaggggcaa aactttttct atcctacttt taacatactt aagtcattgt gttttaaata	5040
catttcttgt agacatcata tagttgggtc tttttaaaaa aatccatact gttgatcttt	5100
gtcttttgat tggatgttt tgaccatttg cagctaattg agttattcat atttgtattt	5160
agatctactg ttttattttc tgtttgtgcc ctctcctttt tattcacatt ctcttctttc	5220
ttttggattg tatctctctc tctctctctc acacacacac acgtatgtga gatatatata	5280
tatctcatg agataatttt atattttttc ttttaacaat tacatatctt ttaaacaact	5340
tgaggagaat agtagcctac tgtattttta cagatgtttt aatttctatt attcttctt	5400
cttttccaac attccatact tccttctggt attctttctc ttctgttctt cagacagaag	5460
ttctgtctga agaactttct taagtatttt ttgtacttgt tttaaagcag ttctgtggc	5520

- 5/24 -

aaaaacttct	tttttccttt	acctgaaaat	attctcattt	caccttcatt	cctaaaggat	5580
attttcactg	catacagaat	tctaggtttg	cagtacattt	agcacataaa	aatattcca	5640
cttccttctg	gottcaatgt	ttctcatggg	aaattggcag	tcttgcaa	aatgtttcc	5700
ctgtaagtaa	tacatcagtt	ttctttgggt	gcatttgatt	tttttttcct	gtgtctttag	5760
tttttagtgc	tttaattatt	acatgtctgg	gcattgattc	ctttgagttt	accctgttta	5820
gggttatctg	agcttcttga	atctataaat	ttataatgtt	agccaaattt	gagaagtgc	5880
taggaattat	ttcttcaaat	acttgctttt	tacacaatag	tctcttctat	gatgatatga	5940
atgtagatc	ttctgatatt	gtccacagg	cccagagag	gctctgttaa	ttttttctcc	6000
taatcttttt	tctctcttgt	ttagattagg	taatttctat	tggcctagcg	acaagttcat	6060
tgactctttc	gttggcttgg	ttcaatctcc	tactctctga	aacacatcca	gatagttttt	6120
atgtgtagt	atatttttca	aagttaaaag	ttccatttga	ttcttcatta	tattttgtat	6180
ttctttggta	agactttcta	tctttccatt	tatttcaagg	gtattcacco	ttaattgtg	6240
aggcatttta	atcatagcta	cgttaaagtc	atctgtgtca	cctaattgtc	ttgtcccact	6300
caagtcaaca	tttttttctc	tacatcttca	tgtgtagagt	aattttgggt	tgaatcctgg	6360
acattttgaa	tatcatgcaa	taaagctttg	ggctctgttt	acttaaagtt	atatggaaaa	6420
tgctgatttt	ctttttttta	ttttagcaac	agttaacct	gttaggatta	ggctgcaaat	6480
tctgactggg	cttctgtggg	ttgtgatttt	aatgtcagtt	ccatttccaa	agccttcgaa	6540
gtaccttcca	tatctatccc	acatgtatac	cactcagtg	ccagtatgag	gtcagagcaa	6600
tactttatcc	tacagtgcag	tttttgaaat	ctgtgataca	ctgttttatg	tcagagtcac	6660
acatgcccag	gggtgacttc	agtagttcat	aaacaacttt	acaaaatcct	ttcccaggct	6720
cctttatctc	tctgatctcc	cttgcacttt	tcagttcctt	ggaattcccc	cgttttggtc	6780
ctttggctag	aaagctttat	cctactctgt	gatttgcttc	ttgtgactgc	tccttcctat	6840
ggggccaagt	gggtgggagga	cagagaaaga	gagagagaga	caaagtaatg	gggatttgcc	6900
ccagtctttt	ggaactacaa	gtcctccaat	cagagggtaa	gattctctcc	ttcatcaaag	6960
ttttgggctc	ctgaggggtc	cccttggtgc	agcgttgct	gttaccacag	tcccaggatg	7020
cctggaggaa	tggaatgcaa	gaggctgata	tgtagagat	nnnnnnnnnn	nnnnnnnnnn	7080
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	7140
nnnnnnnnnn	nnnnnnnnnn	gaattctacc	agatgtacaa	aaaagagctg	gtaccatacc	7200
tactaaaatt	atccccaaaa	ttgaagagga	gggactcctc	cccaactcac	tggcaggcca	7260
gcatcatcct	gataccaaaa	tctgacaaca	gacacaacag	aaaataaaat	ttcaggccaa	7320

- 6/24 -

tatcattaat gaacattgat atanaaatca tcaataaaat actttccagc cgaatccagc	7380
agcacatcat aaaagtaatc caccatgagc aagtaggctt ccttcccaga tgcaaggntg	7440
gttcacatat gcaatcataa atgggtattct cactaagaga ctaagacaaa acaatgatat	7500
ctcatgagnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	7560
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnga attctacagt	7620
atgtattttt gtatatgcat gtgacttcct atgcttttca ttaggctgtg tatagcagta	7680
gtttttcact ttaattgcta tattgtattc tatcattgaa tgtggtacac tttattcatt	7740
ctaattgtca tggatatttg ggttgtttc agtttttgtt acaatgacta atgctgcatt	7800
tattttctaca atatgatagc tgcgatgagt accatatatt ccatttccact ttcctgtagt	7860
atgttgcact gatttttggg agtatgtcct gccttataag gtgtcttate tgtggaaatg	7920
gaacatttat tttattaagt tgtaaataag tttgaaatat caactttcat tacattggga	7980
cagttttctc tgttgacttc cactgaatct tatcatggg atagtgttga tctgagggaa	8040
tgtctaccaa aggtgtccca acaaatattt atcccattaa atacataagt aaattaaaat	8100
tatttttagaa ttgttttgtt actttatata attgataatg tttttcaata tcatgaaatc	8160
agttggaaat cattgtgtc cagtaatgtt gctatttgaa gggctactcc ttctattatg	8220
ttttctcaac ccgaactgaa aagctcattt gtataaaact gcaattggg ggtcaaaatt	8280
tctggccaag gacatgtata tgttaccaat ttgttccaaa gttatggta gagtaatata	8340
gcacttcaat gtatgaaaga aagaaaacac atcaacaaca ctatcaacaa tatacaaaaa	8400
acagttcaac atgtaaaact acctagaacg aatgtgggtg tctcagagtc agtgtttact	8460
gttcgtcttt taaaaatgtt ttccagttct ctggcataat gagagtgatg aaaacaaata	8520
ggcaaaacca aagctcattc tgtttgtgaa aaaaacacta acatgttcac atgttctgaa	8580
aaggaaatat aatttgttca cactcaaaaa tttatgcttt ggagaaatgt aacagctgag	8640
gttgatcatg ataaaacagg aaatgaggaa ggcatgaaaa cactgtaacc cacataagga	8700
cctctttcca aagtgtcaag ctgtttctgt ctggaatttg gaaaagaaat tctttcactc	8760
tagttaatat attctttgcc tcaaagctta actatttttt ctttttcttt ttttgagata	8820
atgactctga acagcatcca actgtacgtt ttactgttgc aaattcacta ttgtaataag	8880
gaatagatgc tctttatttc catagcgtgt aatatggagg aaaacaattt ttgttctatt	8940
tcacctagtc acttgatcta tcaaatcagc acacagaaaa ttaaagcaga tattcaagtg	9000
gtccaatgca aagggttttt ttttaattaaa ataattccaa aaagtaattc ttccaatttt	9060

- 7/24 -

ctgctttgtg	agggtgacca	atztatgaac	tttcacattt	gaaaaacaca	aatgtctcca	9120
gagcggagat	agagttggta	aattctgaca	ctaaatatat	aatgtgacc	aaattatttc	9180
ttggggcctg	tttcctaata	taaaatcaga	ttgtattgat	ctgagttccc	ttctgattta	9240
tcttaaatta	gggaacatta	gtgtaagtca	gatgctctcc	cagagttttc	actaacctgt	9300
tcttggcgct	gttccaaagg	tgctccagag	aagctcctct	gggtagctgc	ctggtggcca	9360
ctgagcaact	ggagattcct	gatgactgca	gctgccaggc	ctgcccagcc	actcccactt	9420
ttgctgacat	gatttatgct	tagcttgggt	agtatagacc	cattccctct	ctttctttct	9480
attacacttc	tttcctccct	tctggcttac	taccaatttt	gggaagaagt	tcagtttttt	9540
tctttggtac	aatgttttac	tgactgtaga	gccaaacttc	acaccttatt	ggaaaaaaaa	9600
ggatagacaa	atagtatttt	tgtgatttta	cagtagctta	aaactatagg	attttttttc	9660
cccattactt	tgcaacaact	gatacttttg	accagttctc	tttcctaagc	atctctctcc	9720
ttgagctatt	acccaaattg	tcctagttct	tctctgcttg	tcctctagct	atagacagtc	9780
actgaggctc	tgtgcttggc	ttctgttcct	ctctcctcat	agggattcct	cctgttggag	9840
ttcatcaagc	tacataggct	gaattatcat	ttttatgact	ctagttgtga	ttactctcta	9900
ggaccacatt	tcctctctgt	aacactttcta	cctgggtact	ctttgagtac	ttcaaaaata	9960
atgtatttaa	ttactgaagt	tgtttttttt	ttgttttttg	ttttgttttg	tttttttgag	10020
atggagtctc	actctgttgc	tcaggctgga	gtgcaatagc	acgatctcag	ctcactgcaa	10080
cctccatgtc	ctgggttcaa	gcaattctcc	tgtcttagcc	tcctgagtag	ctgtgattac	10140
aggcaccgcg	caccacgcct	ggctaatttt	tgtattttta	atagagaagg	ggtttcacca	10200
tgttggccag	gctggctctc	aactcctgat	ctcaggtgat	ccaaccacct	cagcctccca	10260
gagtgtctgg	attacaggcg	tgagccaccg	tgcccagccc	tgaagtaatt	gtttaattaa	10320
aactttctct	tttgtaactt	tctcatttaa	gtatgtgaca	ccattgcttt	tctaagtccc	10380
tagtactgaa	actttatttg	gttttagtcc	accgatctca	ccattcatta	gaaatgtaat	10440
gatttgtcct	gccaaaaagc	ttctgcacat	caaaggaaac	aatcaacaaa	gtgaaaagac	10500
aatccataga	atgagggaaa	atatttgcaa	agtatctggc	aagagcttaa	taaccagaat	10560
atataagcaa	ctctgacaat	tcaatagcaa	aaacaaaaca	aaaaacaaca	aataatctaa	10620
aatggggcaa	aagatctgag	cagacatttc	tcaaaagaag	acttacaaat	ggccaacagg	10680
tatacgaaaa	cattttcaac	atcactaatc	agataaatgc	aatcaaaaac	cacagtgaga	10740
tatctcacc	cagtttgaat	ggcttttata	aaaaagatgg	aacaatagac	ggtagcgagg	10800
atgtggaaaa	aggggaactc	tagtatactg	ttgggtggaa	tgtaaattag	tatagccact	10860

- 8/24 -

aaggaaaaca atagggagtt cttcaaaaaa ctaaaaatag aactaccata tgatccagca 10920
atattattac tgggtatata tccaaaagaa agggaattaa tacattgaag aatatctgta 10980
ttccaatggt tactgcagt ctattcacaa tagccaaaat gtagaatcaa cctaagtgt 11040
gattagtggg tgaatggata aagagaatgt agtatatatg cacaatgaca tattgttcag 11100
ctattaaaaa aaagtgaaat cctgtcattt gaaggaaaat ggaactggag gtcattatgt 11160
taaagtgaat aagccaagcc cagaaagacc aatatcacac gttctccttc atatatggga 11220
gctaaaaaca gaaaaaagtg aatctcatgg agctggagag tagatttgtg gttatcagag 11280
gctgggaatg gtagcgtgga ggagggaatg aagagaggat gattaatgag gacaaatata 11340
cagttagtta ggaggaagac ctagtgtttg atagatcagt atggagacta tagttaacaa 11400
taatctactt gtacatttca atatgtacaa taattcacat gttcacagca taaagaggaa 11460
tgtttaatgt ggtggatatt accctaactt gatatttaca cattatatga atgtatgaaa 11520
atatcaatgt actctgaaaa tatgtccatc tattatgtat taataaaaaa tataagggac 11580
acttactcgt atcaggcaca atatgtgcca atgacatgga atgaaagagc aaataggtga 11640
ctggcactcc ctttcctttc acctcctttc aatacttggc attgagtgga cacatgggtgt 11700
tctagaggct gtgaagcccc cttttcctttt tgtaattga cttccttaga gtctatctca 11760
aaacaacact tcagaagtca tgagggaaaa caagcataat tgtattgact ctttctatat 11820
tttgatatt ctgaataact cttggatgca gaaaacttgt gaaaggcagt ttatttgaaa 11880
ttttccaaag acaaaaagga ataaactgaa tttccttttc ttattaatat agcgttagtt 11940
ctgtcagtct taagcaattt cttttacaaa gtgcaaatta ctgtgaattg agtcctttat 12000
tacaatatat gaggtttgtt ttacagcta agttgagcaa atgttacttt tactttgtaa 12060
agccttggtta tatcgcttag gttcaccagt caacttcatc acattttcca tatactggtg 12120
taatccaaaa taaaatttaa tatgatattc tttgccctgt gtataggtgt aggaaataaa 12180
tattgtttta cataggtaat gaatggttac taagtcaaga cacacttcct tttataaaaa 12240
aatgaaagaa atgtcttatt ttttggtaat aggactttat taaggataaa tttatttaca 12300
gtaaaatact acaaatctta agtgtatgct gcatttaa at tttgacaaat gtatatattt 12360

atgtaaccat catccagatc aagataaaga acatttattt atttatttat ttatttattt 12420
tgagacggag tcttgctctg tcgccaggc tggagtgcag tggcgcgatc tcagctcact 12480
gcaagctcca cccccgggt tcacaccatt ctctgcctc agcctcccga gtagctggga 12540
ctacaggcac ctgccaccac gcccggttaa tttgttgat ttttagtaga gacgggggtt 12600
cacctgttta gccaggatgg tctcgatctc ctgacctcat gatccaccg cctcggcctc 12660

- 9/24 -

ccaaagtgct gggattacag acgtgagcca cgcacccccg gcctaaagaa catttcttta 12720
aattgtctgg tgtcttatca cccattctgc cctgaggcaa gcacagttct gattttcatt 12780
gccatagatg ggtgtcgta ttcagcttca cgtaaattgga agcatactgt acacacttgg 12840
gactggctgc tttggcatac catggcattt ttgagacttt gagttctgtg catcatttta 12900
ttctatttct aattgtgtaa aagtacacct ttgtatgagc ataataccaa tatgacttct 12960
atatgggtcc tagtttaata tagttgattt ctgaatctgc aggagttgtg ctgttggtgt 13020
tgttgttgtt gttcgttttc ttccctttca tcattgacac caggactcta gggttcgggt 13080
gttttgtttg tttgttaatg ccctaggaac aataagaacc tgtcggtttt ccattaagtg 13140
tccatttgat gacttgcaat tttgggaaga ggggaatgtgt tccatggggg ggaatctagg 13200
gatgttttac tatcgcttg tttctagaag gatagtggaa ttgacctgct gaatggccaa 13260
gtggtatatt gccatcttc tgttttaagg actatctgaa caagttccac catggcatcc 13320
attcctgctg ctgttttgat agctgtgatg ataaggtaga tgatccatct agtgactga 13380
tttcttcaact ttactaacct gcttgtctat tctaccccag ccactcaact ccatgagcag 13440
ttctgggcct ttcagagggt ctagcagtgg tgccagcatt gtaatattaa ttctaagaat 13500
cccatacttt gctgtctgcc tctttacatt tcacatgatt tgatgacaca tcatcactgc 13560
tgctactgat tgttttctta ccatgttgag attacctatc catcagctac tgcttactat 13620
ctgctaagcc actggccttc ttgtctatgt gagatatcat ggtctactgc aatactgata 13680
cccttgaaga taccctcaat ttctttgctt tctcttcctc ttccatactc atctggcaaa 13740
tcccaactcc agttatttga gaaaattact caactgaggt gactggtttc actttaaatt 13800
caagatagaa aattcaaata gacaatcaag attgccaagc attcttacct actataaggt 13860
ttgcctttct atttgatgaa ataactgctt tatactttct ttgcttacat gccacatcc 13920
tgcttttcgt agccctcat tccccatagc tgatgacttc acttcattct tcaatgagaa 13980
ggtaaaagcc attggaaaga acaccctcat tattccacca accaatttat aaacttgtca 14040
atatcagtac ctttctctcc ttttcttct ttgtgtagtgg ataaagtctt ctagctttca 14100
aagtcctatt ctctcttttg ctccaggtcc agtcatcacg ttgccagttg aatttagtgg 14160
acccttttct ctctgaactc attttagctc tgagggtcat ttgacacaga ggactacct 14220
gcttttattt tggaagacgc tttttcaatt ccacactctc ctgtatttcc ttccacatca 14280
ctggatgctg cttctcagtc tcctttatgg acttcatatt ctctgtctta tatataaaga 14340
tgtagttcct gcttagagaa ctacaactca ccaaagctt taaatatcat tttctcacgg 14400
tatattggag gccagactag tttaggtcat taagttatat gttcttatta cacatatata 14460

- 10/24 -

ttgtgtttta atttcatgtc atgtatcaca tgtgacaaat tattttgtgt gtactttctt 14520
tgttcaatgt ctttatcctc ttctggattg tctgtagtgc ccatcatgtg tcctacatgc 14580
ctatcacaga gcctggaatg tagcagatgc ccaatatttg cagaatgaat gaattcatat 14640
gtaagtgact gaaagagttc aacaaatggt agttctcttt ccctttctca tctcatattt 14700
acagaagaat ggatacacca gagttatgga gtcagaagat actactatca gttctacaga 14760
agaaagacaa attcatggtg tgatggatgt taccagatgt taaaaattat ggaactgtta 14820
aaaatttttg aactgctcat agatacctcg tacaggcagt tggaaatata gatctagagc 14880
tcagtaaaaa agtcatagtt aaggacctgg atcttcaa atataaaattt tattttttat 14940
atcttttaaa atgtgttgtg ttttgctcct atgtgtttta aaatggagga aaagaggaca 15000
attgggattc aggggtcaaa tgccaggata tgagacttac agtgcactctg gtaaaaatta 15060
aaatgattta aactctgac tgattagaaa aggtcagttc tgaaaaagtt ataaaggaga 15120
ttgtaacaca aatacaattg tattttcttc tctcctccc catagttttt gacaataaat 15180
cctaagcaca gatctgaggc tttttttcct agaaaaaca gaaaaatata cactcaaaca 15240
tagtagtcct ccaaagtcac ggtacatgaa aggttatgag agtttgggga tctgatgaag 15300
ttaattatag tcttacacct ctggataaga cccaaaagca ttcaatttca aacaaaggtc 15360
agggatagtt agaaagtttt atcacacgta agcttgaact caaacaatgt tgcattaaat 15420
agtgcaggtc cagtgagtgt catatttcat agctgtatca gggacttaga cttctttaga 15480
tcttctaaga tcatatcaaa aatactagta tctttttgtt tttaaaaaat aaatgagcat 15540
atgtgtatga gcaaaactga ataaagaagg ctaagattct tgtctcctct gaactatttt 15600
gatctttata agaagctact ctgaggcagc aaggctgttc actgctcggg cactgcgtgt 15660
atcatggggc atttcccctg agctgcagac taacttctgc gaagtactgg gcatgttaac 15720
tttgcaaact cagatttctt tctactaggg tgataaactc atgtttctat gaaataaaca 15780
gtctccaggt tcattagcag atatgtagga gatcattagt attaggagag tcactagaaa 15840
aatcaagcaa gaggagtgga aggaatgtga ggaagaacta agctgaaaat agttttctgt 15900
cctttccatg gctgctggta aatgtagcac tgactctgt ccacacattg aagggtattgt 15960
attaatttca ggctttgagg taaatgaatt gctaattgta aaactgcca tgtttgaaga 16020
tctctttgtc actacttagc ctattttgtc aaaaccaact cctttaactt ttaaagtatt 16080
tttttttctt tacataaagc accaaggcag atgctaattc atttataagc ttttcttctc 16140
aacagtgaat tacagaaaaa tcaatatcta atgggttttg tgcaaagctc agtaataaat 16200

- 11/24 -

agatttcatt ttccattcta aaaaacttgt gccattagga agttgggaga ggaagattac 16260
tagcaattaa acaagaaatg atggtaatgc tacaggcccc ccaaaaagag tatcaagaaa 16320
tcaactatag gtaataaaaa taaactgtta catctgattt agggtagcag gcacaaaaaa 16380
ggtgacagct aatgcatgca acatacctct ctctgattct tcacaatttc acagtctctg 16440
ggcttggtgg gatgtgtacg ttgatgtgtg taaacataat tcagatgagg attatgtttc 16500
agatctattg atctaagata ccccttttcat gaataaccta agatatgaat ttttggggtg 16560
agtaagcata aatgagagga ggttcttttca tatttgagct ttgtggatat aaactcagct 16620
tctccttatt actgttattt tatcgagcta cttgaactat acagataatt tgcttatggc 16680
aagagctagg gattgggaca tattgtccca tataaatatt taaaatatat atttagtatg 16740
gcattgtcac cttggagacc agcagaaata agaagtatat tgcagatgag caaaatttag 16800
tcccagagaat atttgcttag ctcatagct gttaggtaat agtctagcta tgtatatgta 16860
gttgatgca tagcttttga aatgcctact tgaaatatta ctaaatcatt ttgtaataata 16920
atttaaaaa atccctgcct ggaaagtfff atggaatcca tttgtttatg ctccaagcaa 16980
agatatgtaa gtaacaacat caaattagac tcaattcagt gacagttgct atagtttttg 17040
ccaaaatttt ttatttgagc acatgcaaat ttctgaagaa ttacaaatgt gtaatttatt 17100
tacatgactt atttttgata gatatttatt ttttaagtaga aaaatcacao gtctatatat 17160
ggatgaggat aatttataaa cctaatacca taaaatttta cgtcttcaga aatattctca 17220
tctcaagaat atctctttta aaatacatgt aacattttta tacagaataa aatacagaac 17280
gccaaaagct aggttaacttt ttgttctctt ttaaagata atacaatcat ttgctaata 17340
aagtattttt tggaatagtt ttgtcaaaa tagcaacaag cactaatgtt actcattgtt 17400
tattggctga gatcttcaca gaaattgtgt gctaaaaatt atcaggagaa caagggagg 17460
agctgtgtgt ggtattgagg aattctcatt tctgtcatgg gcaagagcaa cagttgctgc 17520
tgctacctat cccagtaga gatacttatt tctctggcag gaagggtgat actcttttcg 17580
ggactgctat gttaaagtgt attattgaca agttcaactg gaaataaaat ttattttcat 17640
tagaaaatac aagtgggtcaa caaaagcttc tgccttttat tgttcccat gttacataa 17700
aacaactttt tttttttttt gagacggggt ottactctat caccagctct ggagtgcagt 17760
ggcttgagct tggctcactg cagctttgac ttccagggt caagtgatcc tcccaactca 17820
gtctcctgaa tagctgaaac tacaggcaca tgcacccaca ccagctgatt ttgcatttt 17880
ttgtagagac ggggttttgc cacattgccc agactgggtc tgaactcctg ggctaaagca 17940
attttttttc cccaccatg ttggcctccc aaagtgagt ctgggattac aggcgtgagc 18000

- 12/24 -

gcttggtgccc agccataaaa tgcctttttaa gtgaaacttt ttttggtgat gtcctttctc 18060
taatttatgt gttcttattt cctttctgtg tcaaaagaga agcacatacc atttaatttt 18120
tactaaagct ttctcagggg cattatttaa agaaatcttt acatataatt aaatattttta 18180
aggtaatgt gtattatttt ctacttttat attgacatta attcctgaca atttgccatg 18240
cttatgatgt atatgtctta gctaagtaat attagctttg tagctcttct ataaaaggta 18300
taataatcta ctcttcaagg tgtcaaatga atctacccta tagatgaaac attttgagat 18360
aattagaatt gttattatta tgatggccaa aaccacaact ccttttgtac caacctaata 18420
tatacataaa atatgcattc tggtcactac tcaccaaata ttgttaaaaa atgaaatttc 18480
tcttttcaaa atacctgcat aatttcctcc aacaagcaca aaaataccat tgagttgaat 18540
ggtagctatt ttacctatag aaggggctca ctgcccctct taagaacttc accaagttca 18600
ctgggttgta ataacacttg cttatctctt tgagttgctt tcagatctag gagctaatag 18660
taggaggagt actgtgtcgt ctgttaaaag taaaaaaaca ttaggtgata gtccacaggg 18720
caaaccagct acctgagcat aggcagaaaa ttgcctccca gttggtattc ataataattaa 18780
tacttccata agcataaagc atacctcttt caggattaag gggtcattgct cttctcatgc 18840
atgggggtga acagcatttc ctcaggattt ggggtggttg ttgagaatgg aaactcgttc 18900
tcatgagcat actgacattg aataagagga catgtagata gaataaacat gaatgtagtt 18960
ccaggagaaa aaattcagac acacagaatt gacaagagcc ttctcaccac caaattgaga 19020
tctagtctga gcaaaaaatg tgagtctgag caaaacaaaa gaaagctaatt attggaaatg 19080
tgcttataac taaaatgtgt ttttcagaac gctgtaaag taagcctatt agagctacac 19140
agaagaccta tttccggaac aattacaact atggtaagga aaatttcatt gttgatgtat 19200
aaactgctgt gttaagagag gtgtgtggat gcacctatga ggatgtgcac atgtaaacga 19260
gtggagccat ttaagaataa ggcttaaaaa cagaaggact attttttagga gcatagtccc 19320
tctgccccga ccaattaaaa tgtctcacac gggcaaaatg cttaacactt ggttgatgtt 19380
gaagcacatc aagcagtatt gtcattctag agccatgcac ttaagtacac agagataagt 19440
atcaagatac atgggactct tccctgcctg gaggtgaag gaggatttga gctcaagttg 19500
ggttgcatag gccatcagtt gtgcttctact gagtgaacag cagagcactt tgatgcataa 19560
ggaactgtta aaagactcaa acttgacttc ttcagtcagt ctatcccatg aaaaacactg 19620
acagatttgg aaagtaacac ttgtagaagt tgctggtatt cagcgaagga aatgtgttgt 19680
tatctagctt gacttaaaat ccgacattct aataaactaa atattaattg aagagccacc 19740
aatccaaatg tagctgacaa acttaactga aggattaaaa agcctgccat ttttgcacat 19800

- 13/24 -

gggactgtgc taagtgattt ccatacatta tctttaattt tctcagctgc cccatgagat 19860
 atattaccat cttgatttta tcaatgatta tcgaacctaa gtagtccctt ctttgattgt 19920
 tcacaatgta ctttgatgc tgggtgtatat tttagagtca taaattcatt tatacagtca 19980
 ttatatattag aaagggcaat taacagagct ttcccgttga cacataaggg actgatattt 20040
 actgtattcc atatgcattt tgattttttc taatatctat ggtattaata aaaggatgt 20100
 tactatctct gcataagact attgatgttg agtcaatgag taaacatgga gattaaacag 20160
 acttttacag tgaaccttat taaaataaga gttttattat tacatttatg tacgctattt 20220
 taaagtatat aggatatata cagcagattt atttcttcta tacttttggg aattgaaagg 20280
 agagaaatga ataagactgt aaaagttagc ttcaatatct tcaaaaatgc agtgtattaa 20340
 .aaaatataaa gttgaaaatt ttgggttttag ttaaactctt ttaaaacaaa tccacctgt 20400
 ttcccaaatt tttcttcaat ggaattaatt tctatgtttt gctgtctact cctcttaaga 20460
 ggattgtagc ttttcttagt caatgatgga tgctctttct aagtttgtaa gcttatgttt 20520
 atgtaagaag tcaaatgcct ggctgcttat ctaagaacct gcttctttac agattgatgt 20580
 gaagaagcct agggaaaagg gagcccttaa tttctaggag attagcttgt taagattgaa 20640
 aaatgctttc tgaactttta tatactgtaa cactaagaat aaaaacatat caattttata 20700
 tcagggaaat gtaattattt atatttaatt tatgtatcct attatcaagt tttattttat 20760
 ctagcttctg tagtgcttca atgggtgttg ctgttatttt tgtaaaaaat tgaatgacaa 20820
 gtcatggaaa gtgaaggctt tagagaaagt agctgaatga atatcataat aaaatgggat 20880
 atggattata attaatggaa aggagcagat aatttctttt ttgtagccaa gtaatagcta 20940
 agttgatcat agtttcagga ttgccatttg tctgggattt cttacagttt ctttggggaa 21000
 gggaggcact caaatggta aatagaagga agaactctga aaaggaagta tattttactt 21060
 ccaaaatatt tttatcttct atatctgcat gtcaagggtg gagcagttgg aaattcagac 21120
 aaaacctcta cttcatttag attaattctg ggaaaaactc cttaagtgtt aaccaaatta 21180
 cttgagtga aaagtaggtc ttgtagaaag atatttattt tggagtgggt acatgtgcta 21240
 aacatttcta atgaaaatag tagacactgg gtaataaat cattagtttg agaacatttg 21300
 cacagttcac taatctctaa tagttgatgg atgctatgaa gcacgttcac ttaagccagg 21360
 acacacaaaa atgcctgcta gctggggttg tgaatttctc atctgattct gccaaatggc 21420
 atttcaaaaa gggcaatctt atcttttggt aaatacagaa tagggagaac atgaatagaa 21480
 agaggctcca cttgccagtc aaaccaagcg aggacagttg ggtagttaca gtttgtttgc 21540
 tacacgttct catacttgag gcaactaatt ctoggaagac tgaaatgtga agctttttaga 21600

- 14/24 -

attccttagt aactaatttc tgtcatgtgt gttgtactgg tagaataagt ttttgtgacg 21660
aaattggaac attaatgggg ggaatggttg attagctggt tttatatttt tcaaggatta 21720
ctttcttcta atcctttata gtccaaaatc tctaaaatat gtgggaaagt actcagatta 21780
aagtggtttg caattatgtg cctaagaagt ttttgaaatg ttgactcatt tggataaacg 21840
gtgttttcct ctatataatg tccatggata gtgaactgga gcaatgaggg tactgggaga 21900
aatgccactg gctttttact gatattttta cttctttctt ggtgttttaa aacttctctt 21960
tttaaaagct catttactag aagcaggaat ctgaacaact tgactctttt aaaggggaga 22020
gatacagaaa agaagcagat ctgtactttc tagattctga gtcacatact ttaactttgc 22080
aactgaaggc aatttgttgt tacattgata agtaaagcaa gtcttgtag atagattggg 22140
tgccaatata tccattgaga gaggtagaat aatgccattg acacgtgaga atatattcca 22200
tagcacgtga gggaaaggaa tgttgagtg tctgacttca atgcagatgg aggaaataag 22260
atctagtggt gaccattttt attcaagaag taaatatagc ccacttagag gagaatataa 22320
aagataatct gtactgaaca caaaggaaaa aaggacctaa tatccgaaca aagaagaaaa 22380
tgtagccac ctaaaggaga atataaaaga taatctgtac tgaccacaaa gaaagaaagg 22440
agctaataac tgaacagtga gccaagatga ggaaggtgaa aagattgcag aaccagtttg 22500
ggaacactac tgagcaggac aattttataa cattacagag gccactact tgagaaaacc 22560
tgttaatgga cctgataaaa cagtgtcgac aaggaataaa gcatatttaa gtattaggtc 22620
tttcatata gtaaaattaa catagttaa attcctcttg gcagcaattg gaactgcaca 22680
tgcatatata cacatccaaa atggaatttt agagaaaaat tgtatcagat tgttgactca 22740
atctttgatt ctctaaatgg atttctcaaa agcacatgca cacacatgca taaatacatg 22800
caaatatata cacagccaca aattggaaag agagcagggc tgagcattag gtcaggtgtg 22860
ttgttttatg tcacacacta cagtctaagg caaaaatac tggctgactt ttctgttcca 22920
ttcgttgaat atcaagtga taagaattaa ctaaagagaa gcctactttg gctttttttg 22980
aatggggtag ctattatgtg tcatattatt atcagctttt cttgtctttt atcccagtca 23040
tttgggctaa agttaagag ataaagacta agtgccatga tgtgactgca gtagtggagg 23100
tgaaggagat tctaaagtcc tctctggtaa acattccacg ggacactgtc aacctetata 23160
ccagctctgg ctgcctctgc cctccactta atgttaatga ggaatatatc atcatgggct 23220
atgaagatga ggaacgttcc aggtaattca ctctttaagg atacagaata actactttgc 23280
ttatcctact ctcatataat ttgtctctag agatgctacc ctgcattttt tacattggga 23340
tctgtctacc ttcttgggga ttacagataa gttttagttg ggtttcttga tttcatatat 23400

- 15/24 -

ggaatgggtg tatgcgtgta gtttataggt ttacatTTTT cttgggagag gggccataac 23460
ttcatttggt tcttaaaggg gtctgtgact tacaaatggg taataattac tgttctaaat 23520
cttagctgtc ctattcaaga ataaacatgt aagtcTTTT tgttcaagta ctgtttcctt 23580
gtacttttca gtaacagggt ttaatttttc aaaaatgttt gcttctttca aaaatacaga 23640
ttactcttgg tggaaggctc tatagctgag aagtggagg atcgacttgg taaaaaagtt 23700
aaggtaagcc tgtattttat gtttgaagta atacacagaa aacaaaacaa aagtagaaaa 23760
ccagccatct gtcattgaaa tgaaatagct gccttcaacc atatgagaaa tattaaaatt 23820
tagtatgcat atatttatat actccaagag ctgaatttgt ctttaatgtc tatttttattt 23880

aaggaatgag ttccctgtat gtgtatcaaa gatctaacta cagttgccct aatgtctctt 23940
cacaaggagt ctagctgtag tccagtcctg gtaggggtgt ttcatgttgt caggctcctt 24000
ctagcttata actctacat ttttaaaggc atcgctttca tactcatgtt cacctggagg 24060
tcacaagatg gctgctctgc ctctagtatc acatgtacat ttcaggcagg aaaaagagga 24120
ggtaaaaggg atttccaaga gcctcatcaa gcagttttta cttagctgca aaacaggcta 24180
gaaaatgtag tcttttttaa cctggggata ttgcttcttt gaacaaactg cagtttgtga 24240
gaaggggaga ataaatgatg ggaaggaaaa taaacaatga tgtccgatga ataaacagca 24300
attcctatgt acaaagtga gctacattca tctgtataga agagatctga ataaattagg 24360
tccttagagc agtaatgtag ccttattttt aagccaaata actcttagat tgtggaattg 24420
atacatTTTT taaaatcaca tagtacagta ctgtattata gaaggttttg tttcttctta 24480
ctgcactaag caaatctgtt ttaaagaaaa gaatgggtg ggggaatgca tggggggaca 24540
aaaaaatctc aaaggatgtg tgggtgggtg gggggagtgg gaactgtacc aaacataata 24600
catcctcttc cttctgcaag tttgaaggct tgggaaacat ttacatggta agttaaacac 24660
ccctgggagg tttatcttat acatctggca cttgacacgg cagacacgtt tgggtttaag 24720
tgacctacaa aaatgtagag ggcattattt tgtaaggaga aactaccctc cagtaagttc 24780
ttctccctt cagtagattt gtcaaaaggg agatgcattt caagttcaga cttgcgggca 24840
ttctctctgc atctctcttc aattagtttg gtgtttatta aagaaaaaag tggctcttgt 24900
tcagtatgtt ggaatctctt actgtcttta atttggtttt ctgcttgcta atgcagaact 24960
ggaatttgct ctaaattcct gtaccagttg gaaaagcttt caatatccct gctgcgaagc 25020
atgtaacaag gcttactgct cctcttttgg taccaaaaca atatcattgt gttgttgaca 25080
ggacacatgg cttgtttgtc ctgtttccca ttgctttctg caatatttcc cattgttttc 25140

- 16/24 -

cgcagggggac ttgcagagtt attattgtaa agtcatttat attctcagtc tggattttaa 25200
tattctaaag gaaagaaacc aataatttga acaattttta attcatatgc agatattgag 25260
ggctaaagag tttgggtaga aaatctcctt tggccagaat tattgagatt cctatttttag 25320
ttttccagtc aatttgaaga tgtgtaaaaa gagttttatg gaggatgtta ttttttcaat 25380
cagatttgct cttagctgag ttttgctctc agtggatgaa tcaataattt tataagaaac 25440
attattatgt ttttttcatg caaatggttg tttacctgag tctcttatag cagacgattc 25500
aagatgtttg gtcttccttg ggaagttgag aatcagactg tttttaaaat atgaagacaa 25560
gattgtggag taaagaatac tgaaatgaag aattaccttc agtaacagaa aaacaataac 25620
acatttacta tgtgctaagc acttttatatt tttattaatc ttcagatgta tctacttata 25680
agtagattct attatcagtt ccaacatata catgatgaaa atgaggtcta gtaagttcaa 25740
gaaacttcat acagtaagtg gtggagttag aatttgaact tgggccatct gacccaagga 25800
tctgatggtg cttgctcttt aaagtgtggt ccctggatca gcaacatagg tatcacctgg 25860
gagcttgta gtaatgaaa tcttcggccc tgtctcagac ctgctaaatg agaatttgca 25920
attttaacaa attccccata tgatatccgt gtttgaagtt tgggaagtac agctgtagtt 25980
cgtattccct acctatctc atgtaatata acttccattt aacttaaatt acatgaaaat 26040
attattgaca ttgttaatat gaaatattac taaacttgat tgtgtttatt aaaaacattg 26100
ctaatacctg ctttgtaaca cttagaacga gggcaggacc ttgtctgttt tattcattgt 26160
acttagtgga tattacttag aaaaaatatt ttgagacccc cccatccctc aaaaaattt 26220
ccattttttt ttcactgctc aaatcacata gactttaaaa ggctatgttc ttatgtgtat 26280
taagagagga tgttgaaaaa tagccttccc ttcaaattct gtgtgtaaatt cttccatttg 26340
ttgagtatct gtgaagcact agcctcctgt atgttacctc aggaaatacc cagagatata 26400
gggagatatt tctgtcatca tttatagaca tggacatttt aagtttaagg tttgataaga 26460
aagcattact aaatacatta tagatgttgc gcttggttaa gtcagagaat aatgttcata 26520
ttgcatcagt tttcttgta ccctgggaaa tagattgata cttttggaaa agtagcaaaa 26580
tgcttcagag atctgaactt gcagtctgac ccagttgtta gaatcatgga aataatgacc 26640
ctgggtgat atgtcttgta ctattttota ttttaaattt cacagcgctg ggatatgaag 26700
cttcgtcatc ttggactcag taaaagtgat tctagcaata gtgattccac tcagagtcag 26760
aagtctggca ggaactcgaa cccccggcaa gcacgcaact aaatcccgaa atacaaaaag 26820
taacacagtg gacttcctat taagacttac ttgcattgct ggactagcaa aggaaaattg 26880
cactattgca catcatattc tattgtttac tataaaaaatc atgtgataac tgattattac 26940

- 17/24 -

ttctgtttct cttttggttt ctgcttctct cttctctcaa cccctttgta atggtttggg 27000
ggcagactct taagtatatt gtgagttttc tatttcacta atcatgagaa aaactgttct 27060
tttgcaataa taataaatta aacatgctgt taccagagcc tctttgctgg agtctccaga 27120
tgttaattta ctttctgcac cccaattggg aatgcaatat tggatgaaaa gagaggtttc 27180
tggatttcac agaaagctag atatgcctta aaacatactc tgccgatcta attacagcct 27240
tatttttgta tgccttttgg gcattctcct catgcttaga aagttccaaa tgtttataaa 27300
ggtaaaatgg cagtttgaag tcaaatgtca cataggcaaa gcaatcaagc accaggaagt 27360
gtttatgagg aaacaacacc caagatgaat tatttttgag actgtcagga agtaaaataa 27420
ataggagctt aagaaagaac attttgcttg attgagaagc acaactgaaa ccagtagccg 27480
ctggggtgtt aatggtagca ttcttctttt ggcaatacat ttgatttggt catgaatata 27540
ttaatcagca ttagagaaat gaattataac tagacatctg ctgttatcac catagttttg 27600
tttaatttgc ttctttttaa ataaacccat tggtgaaagt cttttttttt ctcttctttt 27660
aaaataaatc agaattgccg tattgaccag gaaaagatta tgtatgcacg tgcaccaggg 27720

ttagttttta aaagtacatg gctccataaa aatgctgtag attacagagt gataaaatat 27780
gcagggtttt ttgtttttgt ttttctgtt gtgtgtgtgt ctgtatttgt gtacatgtgt 27840
gtccttgca tccaccccaa ggggtgatta aaatacaggc ctgcaaactg gcctgcactt 27900
tatcatttgg gatttgtgct gcttaatgct cagcgaaaaa tgtctagtaa aatgaattat 27960
ggttgtcagg agagaggtta ttctgacttt tgaaccaatt gcacatttct cattacccaa 28020
gctgtggtga gatcccagg gctgtggtga acaaatgatg cttttataga tgggtccctag 28080
cttacaatga ttcaatttac aatgcttcaa ttaacaatt tgttgactta caatgggttt 28140
atcaggttgt aaccatgc atttcaactt acgatatttt cagtttcgaa tgggtttatc 28200
cagatgtaac cccatcgtaa atggcggagc atctgtactt ccttctctg gcagagttcc 28260
ttagagcctg tgccaattac ttgaaagttt cattctctgc catttacatg tgtgctatta 28320
gtagaaccaa atttctaagc tgtggcgtgc tgaaaataaa atgcttatga agcaaaaatc 28380
gtgctatgtt tcttctaca aggataatca agcaaggatg aaattcattc ctgaacaaaa 28440
gtttcctaaa atggtaaata atcatctatc taaatgttct attttaaaag tgtgagctgg 28500
gtgtggtgag tgcacctgta atccccacta cttgggaggc tgaggaagga agacggctag 28560
agcccatgag gttgaagctg cagtaagcta tgatgcggc cactgtactc cagcgtaggc 28620
gatagagcat gaccctgtct ctgtaacaac acaaaaagtg tgaaagttca ttctacaaat 28680
tggagtcact catatcatac ccaactaaaa tggagttggg aggcctaggg gaaaggcacc 28740

- 18/24 -

cagggtccctg ttccaggaac catttttgca gttcaacaga aaaatcacga agacctaaact 28800
 ctaaccttaa gataaagtta cctagctgct gccactcacc aatcagaact tgctagatcc 28860
 tacaagacgc gcctgctcca gtgactttca ttcaaaacca tntagatcac cttoctctct 28920
 tccccaataa aacccagcc tttctctttg tctntgaaca caactagagg ctacactggg 28980
 ttgtgtgccc agaattacaa ttccaattct tatattccca aataaacctt ttacttggag 29040
 atttatctcc ctatatttaa aggtgacaga gtaagggaaac attcctgcct tctgattgac 29100
 tctccaaagc caacaatttc ccagcccca gaagaaaaca tccccacctt tgcctatagg 29160
 actgcatagt cttttgagct tccaaaagac cacttccaaa gggcttgcca tctgcccatt 29220
 gttgctatat tctcagatgg caaggacatc tcttttgagg tgctgatgtc ctacccttaa 29280
 aagggtgagtt ggggtgtcact ggggagtaat gaagcgccac cgtggatgga ctgcccctga 29340
 gaatctctct tcttggtatg taagcctcaa cttgagcctg aggacagagg ggtggaaagt 29400
 agtgctggtg cctcagcctg ttttctcag gctttctcac gttaactctg taaagaagga 29460
 caagaattga ctacaggaga ggtttaaagg aaccgttatc tcaaattcac aaacctcaac 29520
 tctctcaatg cacagtaagc aatgttaagt aaggaactct tttgatgtat aaaagctgca 29580
 gatgtttcca gcttctgcag ttttttttgg ggggtggggg ttgggttaaag ggggtatgat 29640
 cagtttctgt gtaggaattt gacacacttt atgcttaata taaacaaaac acggccagat 29700
 tcttagattc agcagttttt ttttttttta aaccaccttc cattcggtgc cattttacaa 29760
 cctactgttt taccactacc atctatagtg gcaatgtttg attttgctca cctatatgag 29820
 gcttctgtca gctgttttaa cattttctaat ggtataaagc ccatgaataa agtacatttg 29880
 gtttttcagt ttgatagcta caatatcttc attaataaac tgtgcagacc tcttttggga 29940
 ggatggcgtg atttatcatg atggtcacgt ttctcaggag actgcaaaac catttctact 30000

<210> 2
 <211> 2058
 <212> DNA
 <213> Homo sapiens

<400> 2
 gttgggaaag agcagcctgg gcggcagggg cgggtggctgg agctcggtaa agctcgtggg 60
 accccattgg ggggaatttg tccaaggaag cgggtgattgc cgggggagga gaagctccca 120
 gatccttggtg tccacttgca gcgggggagg cggagacggc ggagcggggc ttttggcgctc 180
 cactgcgcgg ctgcaccctg cccatcctg ccgggatcat ggtctgcggc agcccgggag 240
 ggatgctgct gctgcggggc gggctgcttg ccctggctgc tctctgctg ctccgggtgc 300

- 19/24 -

ccggggctcg ggctgcagcc tgtgagccc tccgcatccc cctgtgcaag tccctgcoct	360
ggaacatgac taagatgccc aaccacctgc accacagcac tcaggccaac gccatcctgg	420
ccatcgagca gttcgaaggt ctgctgggca cccactgcag ccccgatctg ctcttcttcc	480
tctgtgccat gtacgcgccc atctgcacca ttgacttcca gcacgagccc atcaagccct	540
gtaagtctgt gtgcgagcgg gcccggcagg gctgtgagcc catactcatc aagtaccgcc	600
actcgtggcc ggagaacctg gcctgcgagg agctgccagt gtacgacagg ggcgtgtgca	660
tctctccoga ggccatcggt actgcggacg gagctgattt tcctatggat tctagtaacg	720
gaaactgtag aggggcaagc agtgaacgct gtaaagttaa gcctattaga gctacacaga	780
agacctatct ccggaacaat tacaactatg tcattcgggc taaagttaaa gagataaaga	840
ctaagtgcc a tgatgtgact gcagtagtgg aggtgaagga gattctaaag tcctctctgg	900
taaacattcc acgggacact gtcaacctct ataccagctc tggctgcctc tgcctccac	960
ttaatgttaa tgaggaatat atcatcatgg gctatgaaga tgaggaacgt tccagattac	1020
tcttggtgga aggctctata gctgagaagt ggaaggatcg actcggtaaa aaagttaagc	1080
gctgggatat gaagcttcgt catcttgga ctagtaaaag tgattctagc aatagtgatt	1140
ccactcagag tcagaagtct ggcaggaact cgaacccccg gcaagcacgc aactaaatcc	1200
cgaatacaaa aaagtaacac agtggacttc ctattaagac ttacttgcac tgctggacta	1260
gcaaaggaaa attgcactat tgcacatcat attctattgt ttactataaa aatcatgtga	1320
taactgatta ttacttctgt ttctcttttg gtttctgctt ctctcttctc tcaacccctt	1380
tgtaatggtt tgggggcaga ctcttaagta tattgtgagt tttctatttc actaatcatg	1440
agaaaaactg ttcttttgca ataataataa attaaacatg ctgttaccag agcctctttg	1500
ctggagtctc cagatgttaa tttactttct gcacccaat tgggaatgca atattggatg	1560
aaaagagagg tttctggtat tcacagaaag ctagatatgc cttaaaacat actctgcoga	1620
tctaattaca gccttatttt tgtatgcctt ttgggcatte tcctcatgct tagaaagttc	1680
caaagtgtta taaaggtaaa atggcagttt gaagtcaaat gtcacatagg caaagcaatc	1740
aagcaccagg aagtgtttat gaggaacaa caccaagat gaattatttt tgagactgtc	1800
aggaagtaaa ataaatagga gcttaagaaa gaacattttg cctgattgag aagcacaact	1860
gaaaccagta gccgctgggg tggttaatgg agcattcttc ttttggcaat acatttgatt	1920
tgttcatgaa tatattaatc agcattagag aatgaatta taactagaca tctgctgtta	1980
tcaccatagt tttgtttaat ttgcttcctt ttaaataaac ccattggtga aagtcccaaa	2040
aaaaaaaaa aaaaaaaa	2058

- 20/24 -

<210> 3

<211> 23

<212> DNA

<213> Homo sapiens

<400> 3

atagtaggag gagtactgtg tcg

23

<210> 4

<211> 24

<212> DNA

<213> Homo sapiens

<400> 4

aatagtagga ggagtactgt gtca

24

<210> 5

<211> 24

<212> DNA

<213> Homo sapiens

<400> 5

ccctgtggac tatcacctaa tggt

24

<210> 6

<211> 18

<212> DNA

<213> Homo sapiens

<400> 6

ataggccatc agttgtgc

18

<210> 7

<211> 19

<212> DNA

<213> Homo sapiens

<400> 7

cataggccat cagttgtgt

19

<210> 8

<211> 21

<212> DNA

<213> Homo sapiens

<400> 8

tcaaagtgtc ctgctgttca c

21

<210> 9

<211> 22

<212> DNA

<213> Homo sapiens

- 21/24 -

<400> 9
gatcttattt cctccatctg ct 22

<210> 10
<211> 21
<212> DNA
<213> Homo sapiens

<400> 10
atcttatttc ctccatctgc a 21

<210> 11
<211> 20
<212> DNA
<213> Homo sapiens

<400> 11
cgtgagggaa aggaatgttg 20

<210> 12
<211> 22
<212> DNA
<213> Homo sapiens

<400> 12
ttgtctttta tcccagtcac tc 22

<210> 13
<211> 22
<212> DNA
<213> Homo sapiens

<400> 13
ttgtctttta tcccagtcac tt 22

<210> 14
<211> 25
<212> DNA
<213> Homo sapiens

<400> 14
catcatggca cttagtcttt atctc 25

<210> 15
<211> 26
<212> DNA
<213> Homo sapiens

<400> 15
aaaaatgtaa acctataaac tacacg 26

<210> 16

- 22/24 -

<211> 27
<212> DNA
<213> Homo sapiens

<400> 16
gaaaaatgta aacctataaa ctacaca 27

<210> 17
<211> 25
<212> DNA
<213> Homo sapiens

<400> 17
tcttgatttc atatatggaa tgggt 25

<210> 18
<211> 26
<212> DNA
<213> Homo sapiens

<400> 18
acagtacttg aacaagaaag acttat 26

<210> 19
<211> 26
<212> DNA
<213> Homo sapiens

<400> 19
acagtacttg aacaagaaag acttac 26

<210> 20
<211> 28
<212> DNA
<213> Homo sapiens

<400> 20
actgttctaa atcttagctg tcctattc 28

<210> 21
<211> 22
<212> DNA
<213> Homo sapiens

<400> 21
ctcccttttg acaaattctac tg 22

<210> 22
<211> 23
<212> DNA
<213> Homo sapiens

<400> 22
tctccctttt gacaaatcta cta 23

- 23/24 -

<210> 23
<211> 25
<212> DNA
<213> Homo sapiens

<400> 23
gaaactaccc tccagtaagt tcttc

25

<210> 24
<211> 17
<212> DNA
<213> Homo sapiens

<400> 24
ttcgggattt agttgcg

17

<210> 25
<211> 17
<212> DNA
<213> Homo sapiens

<400> 25
ttcgggattt agttgcc

17

<210> 26
<211> 20
<212> DNA
<213> Homo sapiens

<400> 26
gtctggcagg aactcgaacc

20

<210> 27
<211> 18
<212> DNA
<213> Homo sapiens

<400> 27
tgggggcaga ctcttaag

18

<210> 28
<211> 18
<212> DNA
<213> Homo sapiens

<400> 28
tgggggcaga ctcttaaa

18

<210> 29
<211> 27
<212> DNA
<213> Homo sapiens

- 24/24 -

<400> 29

catgattagt gaaatagaaa actcaca

27

<210> 30

<211> 23

<212> DNA

<213> Homo sapiens

<400> 30

gactgaagaa gtcaagtttg agt

23

<210> 31

<211> 22

<212> DNA

<213> Homo sapiens

<400> 31

actgaagaag tcaagtttga gg

22

<210> 32

<211> 21

<212> DNA

<213> Homo sapiens

<400> 32

tgaacagcag agcactttga t

21

<210> 33

<211> 20

<212> DNA

<213> Homo sapiens

<400> 33

gtcggcattc ttatcattca

20

<210> 34

<211> 18

<212> DNA

<213> Homo sapiens

<400> 34

cggcattctt atcattcg

18

<210> 35

<211> 25

<212> DNA

<213> Homo sapiens

<400> 35

aataagtctc atccatactc aaccc

25

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/004178

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used) -

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, EMBASE, CHEM ABS Data, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	CIESIELSKI C.J. ET AL.,: "single nucleotide polymorphism in FZRB in hereditary chondrolysis syndrome" ARTHRITIS AND RHEUMATISM, vol. 46, no. 9, September 2002 (2002-09), page S274, XP009036551 abstract	21-78
A	LOUGHLIN J. ET AL.,: "refined linkage mapping of a hip osteoarthritis susceptibility locus on chromosome 2q" RHEUMATOLOGY, vol. 41, no. 8, August 2002 (2002-08), pages 955-956, XP002298045 the whole document	1-78

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

27 September 2004

Date of mailing of the international search report

11/10/2004

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mueller, F

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/004178

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>LOUGHLIN J. ET AL.,: "functional variants within the secreted frizzled-related protein 3 gene are associated with hip osteoarthritis"</p> <p>PNAS, USA, vol. 101, no. 26, 29 June 2004 (2004-06-29), pages 9757-9762, XP002298046</p> <p>-----</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2004/004178

Box II Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of Item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-20(complete), 41-78 (partial)
polymorphisms in the FRZB gene related to obesity
2. claims: 21-40 (complete), 42-78 (partial)
polymorphisms in the FRZB gene related to osteoporosis